

# Chronic Consumption of Kombucha and Black Tea Prevents Weight Loss in Diabetic Rats

Abbass Morshedi, Mohammad Hossein Dashti-Rahmatabadi\*

Department of Physiology, Shahid Sadoughi University of Medical Sciences, Yazd, Iran

Received: 21 January, 2010 - Accepted: 13 March, 2010

## ABSTRACT

**OBJECTIVE:** Diabetes mellitus (DM) is a chronic metabolic disease which affects many organ systems in the body and is associated with a progressive weight loss. Many diabetic patients used to consume herbal remedies to relieve their symptoms. Black tea and its fermented remedy which is made by cultivating the Kombucha mushroom in a mixture of warm tea and sugar are widely used by diabetic patients throughout the world.

In this study, the effect of chronic consumption of black and Kombucha tea on weight loss was investigated in diabetic Rats.

**MATERIALS AND METHODS:** In this study Streptozotocin (60 mg/kg) was used to induce diabetes mellitus in rats and blood glucose higher than 300 mg/dl was considered as the criterion for the diabetes induction. Control animals received tap water while the sham and test groups consumed sweet black tea solution and different diluents of Kombucha tea solution (25%, 50%, 75% and 100%), respectively for 15 days as their drink. Animals weight was measured before DM induction and on days 1, 3, 7 and 15 after DM induction. Data were analyzed as mean  $\pm$  SEM on different days and in different groups by using T-test and one-way ANOVA.

**RESULTS:** All diabetic animals showed a significant weight loss ( $P < 0.05$ ) which continued progressively in control group, but in animals consuming black or Kombucha tea, there was a progressive weight gain during the experimental procedure, so at the end of the experiment, the animals weight in these groups was the same as their weight before DM induction ( $P > 0.05$ ). At the end of this experiment there was no significant difference in the animals weight in black and Kombucha tea groups ( $P > 0.05$ ).

**CONCLUSION:** According to our findings both black and Kombucha tea induce weigh gain in diabetic rats which may be due to some antidiabetic agents in these remedies.

**KEYWORDS:** Kombucha Tea, Black Tea, Diabetes Mellitus, Weight Loss, Rat.

## INTRODUCTION

Diabetes Mellitus (DM) is a complex disease with a high prevalence. The international diabetic federation estimates that the

prevalence of diabetes is about 7% in the United States (1,2). In Iran more than 2 million adults (7.7%), aged 25-64, suffer from diabetes (3). On the other hand, beside the

---

\*Correspondence: Mohammad Hossein Dashti-Rahmatabadi, Physiology Dept., Shahid Sadoughi University of Medical Sciences, Yazd, Iran. **Tel:** (+98) 351 820 34 10-7. **Email:** dashti-r@ssu.ac.ir


presence of synthetic antidiabetic drugs, a large and increasing number of patients use medicinal herbs or seek the advice of the physicians regarding their use (4-5). One of these herbal medications which are relatively popular in diabetic patients is Kombucha tea (6). The principle element of this beverage is black tea which contains myricetin that mimics insulin in stimulating lipogenesis and glucose transport in rat adipocytes in vitro(7). Therefore, we conducted this study to compare the effect of black tea and Kombucha tea on weight loss in Streptozotocin induced diabetic rats, in a lab trial experimental procedure.

#### MATERIALS AND METHODS


In this study 42 male Wistar Rats weighing  $200 \pm 5$  were divided into six groups. DM was induced by intrapretoneal injection of streptozotocin (60 mg/kg) in all groups. Animals in the first group (control) had free access to tap water, the second group (sham) consumed sweet black tea solution and the remaining 4 groups (test) received different diluents of Kombucha tea (25%, 50%, 75% and 100%) for 15 days instead of tap water. Animals in all groups were weighed before DM induction and on days 1, 3, 7 and 15 after DM induction. The data were analyzed as Mean  $\pm$  SEM of weights on different days and in different groups using T-test and one-way ANOVA.

#### RESULTS


Our results showed that all groups had a significant weight loss due to DM induction ( $P < 0.05$ , Figure 1) which continued progressively in control group (Figure 2), but in sham and test groups there was a progressive weight gain during the 15 days of remedy consumption (Figures 3,4). So there was no significant difference in animals weight in sham and test groups at the end of the experiment and their weight before DM induction ( $P > 0.05$ ). There was also no significant difference in the animals weight ratio (% weight on day 0/weight on day 15) in sham and test groups at the end of this period ( $P > 0.05$ , Figure 5).




**Figure 1-** Effect of Diabetes Mellitus induction on animals weight in different test (K-25-100%), sham and control groups. (n = 7). Weight loss was significant in all groups ( $P < 0.05$ ).




**Figure 2-** The effect of Diabetes Mellitus on weight loss during subsequent days after diabetes induction in control group (n = 7). The weight loss on day 15 as compared with the day 0 was significant ( $P < 0.05$ ).



**Figure 3-** The effect of black tea on weight in Diabetic Rats during subsequent days after Diabetes induction (n = 7). There was no significant difference in weight on day 15 and day 0 ( $P > 0.05$ ).



**Figure 4-** The effect of different concentrations of Kombucha tea on weight in Diabetic Rats during subsequent days after Diabetes induction (n = 7). There was no significant difference in weight on day 15 and day 0 ( $P > 0.05$ ).



**Figure 5-** A comparison between the percent of weight ratio (weight on day 15/weight on day 0) in sham (black tea) and test (100% Kombucha tea) groups 15 days after Diabetes induction ( $n = 7$ ). There was no significant difference in the percent of weight ratio in these groups ( $P > 0.05$ ).

## DISCUSSION

Tea is a beverage with the greatest consumption worldwide. There are 3 categories of tea: black, green, and oolong. Consumption of black tea accounts for 80% of total tea intake (8). Black tea leaves are fermented and contain mostly theaflavins and its major phytochemicals are phenolic compounds like flavonoids and phenolic acids (9). Flavonoids are ubiquitous in plants; almost all plant tissues are able to synthesize flavonoids. The number of flavonoids may be close to 5000. The most important flavonoids are quercetin, kaempferol, myricetin, and chrysin which are found in most fruits and vegetables (10). The content of flavonoids in various beverages like tea and wine is very high (11). Black tea infusions prepared with tea bags (4.0 or 5.0 g) contain 17-25 mg/l quercetin, 13-17 mg/l kaempferol, and approximately 3 mg/l myricetin (8). The contents of catechins range 3-10% (wt/wt solids) (12). Edible plant material also contains numerous weakly estrogenic diphenolic compounds termed phytoestrogens which are lignans, isoflavones, coumestans, and resorcylic acid lactones. The precursors of these compounds are to be found in fiber-rich unrefined grain products, various seeds, cereals, legumes and tea (13-14). Several epidemiological studies suggest that black tea

consumption is associated with a reduced risk of degenerative diseases such as cardiovascular diseases (15), cancers (16) and diabetes mellitus (17). The most effective agents in tea that prevent and/or treat the diseases are flavonoids, content of which is very high in tea (9-10). One of these polyphenolic agents is myricetin which mimics insulin in stimulating lipogenesis and glucose transport in rat adipocytes in vitro. Myricetin-stimulated glucose uptake is possibly due to a change in the intrinsic activity of the glucose transporter, caused by alterations in membrane fluidity or transporter-lipid interactions, as a result of the insertion of myricetin into the membrane bilayer. Thus, myricetin may have a therapeutic potential in management of non-insulin dependent diabetes mellitus by stimulating glucose uptake, without the presence of fully functional insulin receptors (7). Our results also indicate the effectiveness of black tea in reversing weight loss which is one of the most important signs of the streptozotocin induced diabetes mellitus in rats (18). This is despite the fact that tea consumption has a potency of increasing energy expenditure and fat oxidation in normal (non-diabetic) subjects which lead to weight loss (19). On the other hand, there was a similar effect for Kombucha tea which is a beverage made by cultivating a mushroom in black tea. Although the components of Kombucha tea are quite different from that of black tea, antidiabetic effect of this beverage may also be due to polyphenolic residues from the black tea as the basic element of Kombucha tea which needs further investigations.

## ACKNOWLEDGEMENTS

The authors are grateful to Research Vice Chancellor of Shahid Sadoughi University of Medical Sciences for financially supporting this research project.

This article was published in Farsi in Journal of Medicinal Plants Winter 2006; 5(Supplement):42-47.

REFERENCES

1. Inzucchi S, Sherwin R. Chapter 247: Diabetes mellitus. In: Goldman L, Ausiello D. Cecil Textbook of Medicine. 23th ed. Philadelphia, PA: Saunders; 2008: 1727–28.
2. Eisenberg DM, Davis RB, Ettner SL, Appel S, Wilkey S, Van Rompay M, et al. Trends in alternative medicine use in the United States, 1990-1997: results of a follow-up national survey. *Jama* 1998;280(18):1569.
3. Esteghamati A, Gouya MM, Abbasi M, Delavari A, Alikhani S, Alaedini F, et al. Prevalence of diabetes and impaired fasting glucose in the adult population of Iran: National Survey of Risk Factors for Non-Communicable Diseases of Iran. *Diabetes care* 2008; 31(1):96-8.
4. Bennett J, Brown CM. Use of herbal remedies by patients in a health maintenance organization. *J Am Pharm Assoc* 2000;40:353-8.
5. Bent S. Herbal medicine in the United States: review of efficacy, safety, and regulation: grand rounds at University of California, San Francisco Medical Center. *J Gen Intern Med* 2008; 23:854–9.
6. Tietze H. Kombucha: The miracle fungus. Health & Fitness Publication 1997; 109.
7. Ong KC, Khoo HE. Insulinomimetic effects of myricetin on lipogenesis and glucose transport in rat adipocytes but not glucose transporter translocation. *Biochem Pharmacol* 1996;51:423-9.
8. Steele VE, Bagheri D, Balentine DA, Boone CW, Mehta R, Morse MA, et al. Preclinical efficacy studies of green and black tea extracts. *Proceedings of the Society for Experimental Biology and Medicine* 1999;220(4):210-2.
9. Balentine DA, Wiseman SA, Bouwens LCM. The chemistry of tea flavonoids. *Critical Reviews in Food Science and Nutrition* 1997;37(8):693-704.
10. Hollman PCH, Tijburg LBM, Yang CS. Bioavailability of flavonoids from tea. *Critical Reviews in Food Science and Nutrition* 1997;37(8):719-38.
11. Hertog MGL, Hollman PCH, Van de Putte B. Content of potentially anticarcinogenic flavonoids of tea infusions, wines, and fruit juices. *Journal of agricultural and food chemistry* 1993;41(8):1242-6.
12. Bronner WE, Beecher GR. Method for determining the content of catechins in tea infusions by high-performance liquid chromatography. *Journal of Chromatography A* 1998;805(1-2):137-42.
13. Andlauer W, Furst P. Antioxidative power of phytochemicals with special reference to cereals. *Cereal Foods World* 1998;43(5):356-60.
14. Wang H, Murphy PA. Isoflavone content in commercial soybean foods. *Journal of agricultural and food chemistry* 1994;42(8):1666-73.
15. Hertog MG, Sweetnam PM, Fehily AM, Elwood PC, Kromhout D. Antioxidant flavonols and ischemic heart disease in a Welsh population of men: the Caerphilly Study. *American Journal of Clinical Nutrition* 1997;65(5):1489.
16. Lachance PA. Cancer preventive studies: Past, present, and future directions. *Nutrition* 1998;14:237.
17. Day C. Traditional plant treatments for diabetes mellitus: pharmaceutical foods. *British Journal of Nutrition* 1998;80(01):5-6.
18. Carvalho EN, Carvalho NAS, Ferreira LM. Experimental model of induction of diabetes mellitus in rats. *Acta Cirurgica Brasileira* 2003;18:60-4.
19. St-Onge MP. Dietary fats, teas, dairy, and nuts: potential functional foods for weight control? *American Journal of Clinical Nutrition* 2005;81(1):7-15.