

## Effect of an Aerobic Exercise Course on PI3K and AKT1 Expression and Neural Muscle Insulin Resistance in Diabetic Rats

Mahdi Nadi<sup>1\*</sup>, Abdolali Banaeifar<sup>2</sup>, Sajad Arshadi<sup>3</sup>

1. PHD Student, Islamic Azad University of Tehran South, Tehran, Iran.  
2. Associate Professor, Islamic Azad University of Tehran South, Tehran, Iran.  
3. Assistant Professor, Islamic Azad University of Tehran South, Tehran, Iran.

### \*Correspondence:

Mahdi Nadi, PHD Student, Islamic Azad University of Tehran South, Tehran, Iran.

Tel: (98) 913 373 1507

Email: mahdinadi61@gmail.com

ORCID ID: (0000-0003-1837-9373)

**Received:** 15 February 2021

**Accepted:** 14 June 2021

**Published in September 2021**

### Abstract

**Objective:** The purpose of the present study was to investigate eight weeks of aerobic workout on PI3K and AKT expression as well as insulin resistance (IR) of muscle in diabetic rats by nicotinamide - streptozotocin.

**Materials and Methods:** This laboratory study was conducted on 14 male Wistar rats (8 to 10 weeks) with a weight range of 201 to 250 g and induction of type 2 diabetes (one week). These mice were classified into 2 groups: aerobic training and control group. No exercise was given to the control group during the study, while the aerobic exercise program was run for 5 weeks a week with a gradual increase of speed (10 to 25 m / min) and time (15 to 40 minutes) in running treadmill was performed for the aerobic training group. Assay of gene expression in both study groups was by rt-PCR. For statistical analysis, the SPSS 19 software was used. The variables were compared between the two groups using one-way ANOVA.

**Results:** The findings showed that there was no significant difference between the two groups in terms of IR in diabetic rats ( $3.85 (\pm 0.39)$  vs.  $5.26 (\pm 0.55)$ ;  $P$ -value= 0.345). The expression of AKT ( $2.37 (\pm 2.33)$  vs.  $1.000 (\pm 0.001)$ ;  $P$ -value= 0.042) and PI3K ( $2.87 (\pm 2.54)$  vs.  $1.000 (\pm 0.001)$ ;  $P$ -value= 0.028) in the neural muscle of the training group compared with control group had a significant increase.

**Conclusion:** It seems that performing eight weeks of aerobic exercise could be a strong stimulus for PI3K and AKT gene expression.

**Keywords:** Exercise, AKT, PIK3R3, Insulin resistance, Diabetes mellitus

### Introduction

Type 2 diabetes mellitus (T2DM) is a complicated metabolic disorder that influences various organs and is characterized by hyperglycemia (1). Nowadays, there is no persistent and effective treatment for this disease; therefore, the

mechanisms that may display a role in the progression of diabetes and executable ways for treating this disease should be identified (2).

Protein kinase B (AKT) is a group of serine or threonine kinases that play a vital role in

signaling pathway of phosphoinositide 3-kinase (PI3K) (3). According to studies, balance in the PI3K/ AKT signaling pathway is essential for normal metabolism, and damage to this pathway leads to increased obesity, IR, and T2DM. The IR, in turn, exacerbates damage in the PI3K/ AKT pathway and makes a destructive cycle (4).

Aerobic exercise is the best practice model for interacting with diabetics who can improve many of the side effects associated with the disease (5). Evidence suggests that the mechanism of action of aerobic exercise on glucose homeostasis and insulin function mainly depends on muscle function (6). Muscle makes up about half of your body weight and the main place of consumption is glucose (7). Muscle contraction that occurs as a result of aerobic exercise has an insulin-like role, it causes a lot of glucose to enter the cell and produce energy (7).

Given that the highest amount of glucose in the bloodstream is removed by skeletal muscle (7), and since the effects of diabetes and exercise, especially aerobic workout, on the activation of the PI3K/ AKT pathway in the muscles of diabetics have not yet been determined. Hence, this study discusses the role of PI3K/ AKT signaling in skeletal muscle, adipose tissue, liver, brain, and pancreas. It also provides evidence of factors affecting the PI3K/ AKT pathway that may be effective in treating obesity and T2DM.

## Materials and Methods

This laboratory study was performed on 30 male Wistar rats (age: 10 weeks / mean weight  $230 \pm 10$  g) who were randomly divided into two groups (control, aerobic). In order to adapt the animal to the laboratory environment, rats were initially maintained in the laboratory for 2 weeks without intervention. During the first sessions and exercises in the experimental groups, we encountered attrition; Eight mice in

each group died during the study. Therefore, the final sample size of 7 male rats in each group was determined. After separating the rats into diabetic and healthy groups, T2DM was induced in the intra-peritoneal injection of nicotinamide and Streptozotocin (STZ) in diabetic groups.

The group included 7 male 10-week old male Wistar rats who, after familiarizing themselves with the lab environment and inducing T2DM, participated in an aerobic training course from the 12th week. The exercise program was conducted for 8 weeks in 5 sessions per week with a gradual increase in speed (10 to 25 m / min) and time (15 to 40 minutes) in the form of running on the treadmill. All mice were described 48 hours after the last workout. The diabetic control group consists of 7 male 10-week old male Wistar rats with T2DM.

There was a reverse primer in the kit. But the Forward primer is designed with iodine. In fact, the forward primer is the same as the adult microRNA sequence, but it should be checked for the melting temperature ( $T_m$ ), so that if its melting temperature is not matched to the reverse primer, changes to its structure are given. After designing the primer by a geneticist using Primer 3 software, the order was made to make the company a pioneer and was prepared after a week. In addition, the RNA-polymerz2 gene was used as control gene. Table 2 shows the pattern of primers.

## RNA extraction

The RNA was extracted from the neural tissue of Rneasy protect mini kit (QIAGEN) according to the company's instructions. So, we scraped 20 milligrams of tissue from scalpel into microchips, and then the RNA was extracted using the RNeasy Protect kit in accordance with the instructions of the German manufacturer.

To extract RNA, 20 mg of crushed tissue was inserted into the microtype and The RNA was

**Table 1. Aerobic exercise protocol**

Week	First	Second	Third	Fourth	Fifth	Sixth	Seventh	Eighth
Speed (m/min)	10	10	15	15	20	20	25	25
Duration (min)	15	15	20	20	20	20	40	40

then extracted using the RNeasy Protect mini kit (QIAGEN) according to the manufacturer's instructions. To determine IGF1 mRNA, AKT1 mRNA and PI3K mRNA by RT-Real time PCR, Rotorgen 6000 system was used using the One Step SYBR TAKARA single-step kit according to the manufacturer's instructions. Thermal cycle protocol used by Rotogen device in Real time-PCR includes: 42° for 20 minutes, 95° for 2 minutes, 40 cycles at 94° for 10 seconds and was 60° for 40 seconds. RNA Polymrase II was used as a control gene to determine the expression of the studied genes. CT of the reactions was extracted and recorded by Real time-PCR software. A comparative  $\Delta\Delta CT$  method was used to quantify the expression of mRNA.

### Statistical analysis

The Kolmogorov-Smirnov test was used to ensure the normal distribution of data. Descriptive statistics were used to describe the data and draw charts. One-way ANOVA was used to evaluate the variables between diabetic groups (control, aerobic), and LSD test was used to perform complementary tests if needed. The significance level was considered as  $< 0.05$ . Statistical analyzes were performed using SPSS/ Win software version 19.

### Ethical considerations

The protocol of this study was approved by the Ethics Committee of the Institute of Physical Education and Sports Sciences (IR.SSRC.REC.1398.047).

### Results

Information on body weight, IR, AKT, and PI3K was reported in table 3. Comparison of body weight values in the duodenal axis of diabetic rats demonstrated a considerable increase in the endurance workout group before and after the intervention ( $P$ -value= 0.006). Also, at the end of the study, body weight showed a remarkable difference between the two groups ( $P$ -value= 0.001). There was no statistically significant difference between the IR (Fasting Insulin  $> 60$  pmol/l) in the intervention and control groups ( $P$ -value= 0.345). In another word, endurance training could not significantly reduce IR in diabetic rats. This trial demonstrated that compared to the control group, AKT expression ( $P$ -value= 0.042) and PI3K expression ( $P$ -value= 0.028) were significantly increased in the duodenal muscle of diabetic rats in the intervention group. The changes of these variables were doubled in the endurance-training group (Table 3).

### Discussion

Findings from this study indicated that there is

**Table 2. Primer pattern used in research.**

Genes	Primer sequence	Product size	T m	Gene Bank
AKT1	For: AGGAGGTCATCGTTGCCAAG Rev: GCTCACGAGACAGGTGGAAG	159 bp	60	NM_001191052.1
PI3K	For: ACTGAGATGGAGACACGGAAC Rev: GCATCCAAGGGTCCAGTTAGTG	159 bp	60	NM_001191052.1
RNA polymraseII	For: ACTTTGATGACGTGGAGGAGGAC Rev: GTTGGCCTGCGGTCGTTC	164 bp	60	NM_008759265.1

**Table 3. The comparison of mean ( $\pm$  standard deviations) of variables**

Variables	Group	Baseline	After 6 weeks	$P$ -value*	$P$ -value**
Weight	Endurance training	230.85 ( $\pm$ 2.47)	284.85 ( $\pm$ 10.77)	0.006	
	control	230.42 ( $\pm$ 2.82)	292.25 ( $\pm$ 8.05)	0.104	0.001
IR	workout	-	3.85 ( $\pm$ 0.39)	-	
	control	-	5.26 ( $\pm$ 0.55)	-	0.345
AKT	workout	-	2.37 ( $\pm$ 2.33)	-	
	control	-	1.000 ( $\pm$ 0.000)	-	0.042
PI3K	workout	-	2.87 ( $\pm$ 2.54)	-	
	control	-	1.000 ( $\pm$ 0.001)	-	0.028

\* $P$ -value: within groups. \*\*  $P$ -value: between groups.  $P$ -value was obtained from Independent samples T-test (2-tailed).

Abbreviations: IR: Insulin Resistance; AKT: Calorie restriction; PI3K: Body Mass Index.

a significant difference after 8 weeks of aerobic exercise activity between the control group and aerobic training in body weight changes and expression of AKT1 and PI3K1. IR due to T2DM leads to impaired function of beta cells (8). The results of many studies show that high-intensity exercise is a very important factor in improving insulin levels (9); therefore, endurance exercise can improve muscle performance, workout capacity, and ultimately prevent disease and early mortality (10).

Results of our study is in line with the results of some studies (11). Kazior et al. expressed that seven weeks of endurance-resistance training significantly increased oxygen uptake and levels of AKT (12). Chavanelle et al. demonstrated a considerable increase in the expression of AKT protein in the bipartite and fetal muscle by continuous high-intensity exercise. Another study reported that exercise activates AKT and reduces oxidative stress and thus reduces apoptosis in neurons (13). Chih-Hsueh et al. displayed that physical activity and resveratrol supplementation lead to increased activity of the PI3K / AKT signaling pathway (14).

While the results obtained from some other studies are inconsistent with the findings of our research. Sherafati Moghadam et al. in a study showed that four weeks of HIIT in neural muscle of T2DM rats did not cause a significant difference in AKT1 protein levels (15). Mascher et al. conducted a study to assess the rate of protein synthesis deficit in the early recovery period after a severe aerobic workout in the lack of dietary supplements. The findings of this study showed that stimulation of the mTOR signaling pathway had no significant effect on AKT phosphorylation (16). Shadmehri et al. stated that there was no significant difference in the 8-week endurance training in AKT1 protein content (17).

Clinical studies have shown that PI3K pathway activity is reduced in the skeletal muscle of T2DM patients (18). Exercise activates AMPK, increases insulin receptors

substrate 1 and 2 (IRS1 and IRS2) and PI3K/ AKT in skeletal muscle (19); Thus, exercise increases the expression of the GLUT4 gene by activating the AMPK pathway and improves glucose uptake into muscle cells (20, 21). Reinforcement of the PI3K/ AKT pathway and BCL-2 family also prevents diabetes-induced apoptosis (22). On the other hand, due to the effect of T2DM on protein homeostasis and impaired protein synthesis, physical activity leads to improved IR, T2DM, and ultimately improved protein synthesis (23).

This study has some limitations. Small sample size, short median time, and non-consideration of calorie intake by rats are among the limitations of the present study, which is suggested to be examined in future research. In addition, the use of alternative pathways to regulate glucose metabolism can determine the intracellular defect in the insulin signal cascade and thus help in the treatment of obesity and T2DM.

## Conclusions

Co-activity is a strong stimulus that can make changes in signal transduction and cellular metabolism, which varies with intensity, type, and duration of exercise. Therefore, the selection of sports activities with different conditions (severity, type, and duration) is important for biochemical and morphological adaptation in skeletal muscle.

Endurance and aerobic exercises, by making their biochemical changes in the muscles, increasing capillary density, and increasing oxidative enzymes, can improve the transport and metabolism of glucose and increase the ability of insulin binding to muscle cell receptors, and in the result is a reduction in the need for insulin. Therefore, it seems that the implementation of eight weeks of aerobic exercise can be a strong stimulant for the expression of PI3K and AKT gene in neural muscle in T2DM rats.

## Acknowledgments

The authors thank all those who contributed to this research. We also thank the Islamic Azad University, South Tehran Branch for its financial support.

## Funding

This study did not receive any specific grant from funding agencies.

## Conflict of Interest

None.

## References

1. Rask-Madsen C, King GL. Vascular complications of diabetes: mechanisms of injury and protective factors. *Cell metabolism*. 2013;17(1):20-33.
2. Huang X, Liu G, Guo J, Su Z. The PI3K/AKT pathway in obesity and type 2 diabetes. *International journal of biological sciences*. 2018;14(11):1483.
3. Shiojima I, Walsh K. Regulation of cardiac growth and coronary angiogenesis by the Akt/PKB signaling pathway. *Genes & development*. 2006;20(24):3347-65.
4. Cai S, Khoo J, Channon KM. Augmented BH4 by gene transfer restores nitric oxide synthase function in hyperglycemic human endothelial cells. *Cardiovascular research*. 2005;65(4):823-31.
5. Yang Z, Scott CA, Mao C, Tang J, Farmer AJ. Resistance exercise versus aerobic exercise for type 2 diabetes: a systematic review and meta-analysis. *Sports medicine*. 2014;44(4):487-99.
6. Azari N, Rahmati M, Fathi M. The Effects Of Endurance Exercise On Blood Glucose, Insulin And Insulin Resistance In Patients With Type II Diabetes: A Systematic Review And Meta-Analysis Of Studies In Iran. *Iranian Journal Of Diabetes And Metabolism*. 2018;17(2):65-78.
7. Ivy JL. Role of exercise training in the prevention and treatment of insulin resistance and non-insulin-dependent diabetes mellitus. *Sports medicine*. 1997;24(5):321-36.
8. Suhara T, Baba Y, Shimada BK, Higa JK, Matsui T. The mTOR signaling pathway in myocardial dysfunction in type 2 diabetes mellitus. *Current diabetes reports*. 2017;17(6):38.
9. Marliss EB, Vranic M. Intense exercise has unique effects on both insulin release and its roles in glucoregulation: implications for diabetes. *Diabetes*. 2002;51(1):S271-83.
10. Crane JD, MacNeil LG, Tarnopolsky MA. Long-term aerobic exercise is associated with greater muscle strength throughout the life span. *Journals of Gerontology Series A: Biomedical Sciences and Medical Sciences*. 2013;68(6):631-8.
11. Li J, Liu Y, Liu B, Li F, Hu J, Wang Q, et al. Mechanisms of aerobic exercise upregulating the expression of hippocampal synaptic plasticity-associated proteins in diabetic rats. *Neural plasticity*. 2019;2019.
12. Kazior Z, Willis SJ, Moberg M, Apro W, Calbet JA, Holmberg HC, et al. Endurance exercise enhances the effect of strength training on muscle fiber size and protein expression of Akt and mTOR. *PloS one*. 2016;11(2):e0149082.
13. Chavanelle V, Boisseau N, Otero YF, Combaret L, Dardevet D, Montaurier C, et al. Effects of high-intensity interval training and moderate-intensity continuous training on glycaemic control and skeletal muscle mitochondrial function in db/db mice. *Scientific reports*. 2017;7(1):1-10.
14. Lin CH, Lin CC, Ting WJ, Pai PY, Kuo CH, Ho TJ, et al. Resveratrol enhanced FOXO3 phosphorylation via synergetic activation of SIRT1 and PI3K/Akt signaling to improve the effects of exercise in elderly rat hearts. *Age*. 2014;36(5):1-0.
15. SherafatiMoghadam M, Salesi M, Daryanoosh F, HematiNafar M, Fallahi A. The effect of 4 weeks of high intensity interval training on the content of AKT1, mTOR, P70S6K1 and 4E-BP1 in soleus skeletal muscle of rats with type 2 diabetes: An experimental study. *Journal of Rafsanjan University of Medical Sciences*. 2018;17(9):843-54. (in Persian)
16. Mascher H, Ekblom B, Rooyackers O, Blomstrand E. Enhanced rates of muscle protein synthesis and elevated mTOR signalling following endurance exercise in human subjects. *Acta Physiologica*. 2011;202(2):175-84.
17. Shadmehri S, Moghadam MS, Daryanoosh F, Golbar SJ, Tanideh N. The effect of 8 weeks endurance exercise on the content of total and phosphorylated AKT1, mTOR, P70S6K1 and 4E-BP1 in skeletal muscle FHL of rats with type 2 diabetes. *Journal of Shahid Sadoughi University of Medical Sciences*. 2019;26(12):1063-74. (in Persian)
18. Zhang BB, Zhou G, Li C. AMPK: an emerging drug target for diabetes and the metabolic syndrome. *Cell metabolism*. 2009;9(5):407-16.
19. Christ CY, Hunt D, Hancock J, Garcia-Macedo R, Mandarino LJ, Ivy JL. Exercise training improves muscle insulin resistance but not insulin receptor

- signaling in obese Zucker rats. *Journal of Applied Physiology*. 2002;92(2):736-44.
20. Banks EA, Brozinick Jr JT, Yaspelkis III BB, Kang HY, Ivy JL. Muscle glucose transport, GLUT-4 content, and degree of exercise training in obese Zucker rats. *American Journal of Physiology-Endocrinology And Metabolism*. 1992;263(5):E1015-20.
21. Brozinick Jr JT, Etgen Jr GJ, Yaspelkis 3rd BB, Kang HY, Ivy JL. Effects of exercise training on muscle GLUT-4 protein content and translocation in obese Zucker rats. *American Journal of Physiology-Endocrinology And Metabolism*. 1993;265(3):E419-27.
22. Radak Z, Kumagai S, Nakamoto H, Goto S. 8-Oxoguanosine and uracil repair of nuclear and mitochondrial DNA in red and white skeletal muscle of exercise-trained old rats. *Journal of Applied Physiology*. 2007 Apr;102(4):1696-701..
23. Gougeon R, Morais JA, Chevalier S, Pereira S, Lamarche M, Marliss EB. Determinants of whole-body protein metabolism in subjects with and without type 2 diabetes. *Diabetes care*. 2008;31(1):128-33.