Evaluation of Immunoglobulin A in Diabetic Patients and its Relation with Oral Complications

Sodabeh Farahnak¹, Robab Sheikhpour^{2,3*}, Foad Iranmanesh⁴

1- Department of Endodontrics, School of Dentistry, Shahid Sadoughi University of Medical Science, Yazd, Iran

2- Department of Physical Activity and Sport Science, Taft Branch, Islamic Azad University, Taft, Iran

3-Hematology and Oncology Research Center, Shahid Sadoughi University of Medical Science, Yazd, Iran

4- Department of Endodontrics, Dental School , Rafsanjan University of Medical Science, Rafsanjan, Iran

*Correspondence:

Robab Sheikhpour, Department of Physical Activity and Sport Science, Taft Branch, Islamic Azad University, Taft, Iran

Hematology and Oncology Research Center, Shahid Sadoughi University of Medical Science, Yazd, Iran

Email: robab.sheikhpour@iauyazd.ac.ir Tel: (98) 913 152 2462

Received: 15 May 2015

Accepted: 05 Jun 2016

Published in November 2016

Abstract

Diabetes mellitus is a group of metabolic diseases caused by a combination of insulin resistance and impaired insulin secretion by pancreatic β cell. In 2007, 246 million people (roughly 6%) were affected by diabetes worldwide and it is estimated that this will increase to 380 million in 2025. Diabetes is associated with several long-term complications such as cardiovascular disease, nephropathy, retinopathy, neuropathy and complications. In addition, Diabetes mellitus causes an increased risk of morbidity because of infection disease. It seems that, the frequency of infections associated increased Immunoglobulin-A (IgA) deficiency in these patients. Therefore a low IgA secretion rate is suspected to be one of these mechanisms. Moreover, it shows a main antiviral activity by neutralizing toxins and viruses. It by inhibiting the attachment replication of pathogenic microorganisms prevents colonization of these pathogens. Therefore, it acts as a first line of defense against pathogens and early detection immunoglobulin A deficiency in diabetic patients can prevent the vicious cycle of recurrent infections and reduces risk for morbidity and metabolic decompensation. Moreover, the Salivary-IgA is the widespread immunoglobulin in mixed saliva and is assumed to be an important factor for adaptive immunity in the oral cavity. Therefore, according to these studies, Immunoglobulin A, its mechanism, IgA deficiency and diabetes and its relation with oral complications are explained in this paper.

Keywords: Immunoglobulin A, Diabetic patients, Immunodeficiency, Oral complications

Introduction

iabetes mellitus is a group of metabolic diseases caused by a combination of insulin resistance and impaired insulin secretion by pancreatic β cell.

Today, more than 200 million people in worldwide have type 2 diabetes. The total number of people with diabetes is expected to reach 380 million worldwide in 2025 (1-4).

Diabetes is associated with several long-term complications such as cardiovascular disease, nephropathy, retinopathy, neuropathy (5), increased risk of morbidity from infectious diseases (6,7) and some oral complications xerostomia, tooth loss, gingivitis, periodontitis, odontogenic abscesses and soft tissue lesions of the tongue and oral mucosa (8). Moreover, alterations in salivary flow rate compositions and can affect development, symptoms and severity of oral changes in diabetic patients. On the other hand, higher rate of dental problems in diabetic patients is associated with salivary dysfunction, existing acidogenic microorganisms, poor glycemic control, poor dental hygiene and higher dental plaques (8). In addition, Diabetes mellitus causes an increased risk of morbidity because of infection disease, like as pneumonia and urinary tract infections. Several studies reported that diabetes mellitus is associated with impaired chemotaxis of the immune cells, defective phagocytosis of macrophages, and increased production of free radicals (9). Several mechanisms can cause susceptibility of infections in patients with diabetes (9). Immunoglobulin A and G via affecting cavity microorganisms in the saliva, gingival and plasma prevent bacterial metabolism and adhesion of microorganisms to the oral tissue (10). It seems that the increased frequency of infections associated with IgA deficiency in these patients (11). Therefore a low s-IgA secretion rate is suspected to be one of these mechanisms. However, these patients are more susceptible to frequent infections, autoimmune disorders, gastrointestinal diseases and atopy (11). Branco-de-Almeida et al reported that DM patients were associated with lower mean levels of s-IgA than non-DM patients, while Ya vuzyilmaz et al, reported that DM patients had higher mean levels of s-IgA than healthy controls (9). The aim of this study was to evaluate serum and salivary IgA level in another patients. In immunoglobulin A (IgA) and its mechanism are explained. Immunoglobulin A deficiency

(IgA deficiency) and autoimmune disorders are discussed in next section. In osection 2, 3, Immunoglobulin A deficiency (IgA deficiency) and Diabetes and salivary immunoglobulin A and oral complication are explained.

Immunoglobulin A (IgA) and its mechanism

Immunoglobulin- A (IgA) is the second most abundant immunoglobulin (12) in the body and two forms can be identified: IgA1 and IgA2. Immunoglobulin A1 found mainly in the serum in monomeric form (13) and cannot be transported to mucous secretions (9). IgA2 (dimeric) is found in exocrine secretions and mucous defense systems like saliva, tears, colostrum and nasal and gastrointestinal secretions (14,15), bronchoalveolar lavage fluid, urine (9) and plays a main role of antigen excretion (9). The regulation of secretion and synthesis of secretory IgA is dependent on antigenic stimulation and strong neuroendocrine function like stress, exercise, pregnancy and menstrual cycle (11). In addition, it can protect mucous membranes against infections via activating the alternative way of the complement and prevent binding of virus to epithelial cells of the respiratory, gastrointestinal and urogenital Therefore, it shows a main antiviral activity (14) by neutralizing toxins and viruses (9). It by inhibiting the attachment and replication of pathogenic microorganisms colonization of these pathogens (9), and may be protective against periodontal disease (16). Therefore, it can act as a first line of defense against pathogens (16). Study of mucosal humoral immunodeficiency in humans showed that the absence of secretory IgA results to increase mucosal infections and respiratory tract infections. In addition suppression of s-IgA causes increased incidence of upper respiratory tract infection (9). Studies showed that IgA proteases can cleavage IgA in vivo, resulting in generation of intact Fab alpha and (Fc alpha) 2 fragment (17). When bacteria are exposed to Fab alpha fragments, it released

from IgA after cleavage by IgA protease, their surface antigens are likely to be occupied by Fab alpha fragments. These Fab alpha fragments left on the bacterial surface may mediate adhesion. Together, these results indicate that IgA proteases, by promoting adherence, contribute the pathogenic potential of bacteria in the oral (17). Early detection of immunoglobulin A deficiency in diabetic patients can prevent the vicious cycle of recurrent infections and reduces risk for morbidity and metabolic decompensation (18).

• Immunoglobulin A deficiency (IgA deficiency) and autoimmune disorders

IgA deficiency is the commonest type of immunodeficiency (19-24) and associated with autoimmune disorders (14). Prevalence of immunoglobulin A deficiency (IgA-D) as most common primary immunodeficiency disease (25,26) varies widely within different geographical regions from 1/143 to 1/185000 (11,18). Immunoglobulin A deficiency is associated with autoimmune disease such as systemic lupus erythematus, Graves' disease, celiac, recurrent parotiditis, inflammatory bowel syndrome, Crohn's disease, juvenile idiopathic arthritis and type 1 diabetes (11).

Immunoglobulin A deficiency (IgA deficiency) and Diabetes

The association between type 1 diabetes and IgA-D has long been recognized in many populations (18). High prevalence of IgA-D has been found in children and adults (18). Prevalence of autoimmune thyroid was significantly higher in IgA-D patients (18). Moreover, the prevalence of IgA deficiency in type 1 diabetes is estimated between 0.4 % and 5.4%. It is more than 10 times the prevalence of the general population (18). Moreover, prevalence of Immunoglobulin A deficiency in Iranian patients with diabetes mellitus reach to 0.7% (1:150), which is much higher than other prevalence in general populations (27). Greco et al, reported in diabetic patients, because of the availability of serological screening for celiac disease (CD), more cases of IgA-D are now being diagnosed (18). Mohiti-Ardekani et

al. reported that secretory IgA levels in Iranian diabetic patients were higher than in nondiabetics patients (28). Another performed in Western part of Sicily (Italy) and reported that IgA-D was seen in eight out of the 150 subjects (children and adult patients) with type 1 diabetes. They reported that Immunoglobulin- A deficiency was found in 5.3% of patients (18). In one study, no significant differences were found in salivary IgA levels between diabetic patients and control group (6). In another study, the mean level of IgA concentration is low in diabetic patients than control, it seems that infection and other autoimmune disorders are absent in these patients (11). Akefeh et al, reported that there was no significant association between serum and salivary IgA level among diabetic and nondiabetics and also, no association between serum and salivary IgA levels (11). Yavuzyilmaz et al, demonstrated a significant increase in salivary IgA levels in diabetic patients. Vaziri et al, reported that there was no difference in salivary IgA between type 1 and 2 diabetic patients and control group. Also, lower salivary flow rate was detected in diabetic patients than control group (6). Studies reported that these differences can be due to diversity in sample selection criteria, study design (6), or detection methods (10). Moreover, these differences may be due to differences in the type of saliva collected (stimulated or unstimulated), the salivary collection methods, the stage of the disease, and the metabolic control status of the disease (6). Sayarifard in another study reported increased prevalence of immunoglobulin (Ig) A deficiency in a number of autoimmune diseases and type 1 diabetes mellitus (DM1) (27). Smith Jr et al, reported that there is an increased IgA deficiency prevalence in juvenile-onset insulin-dependent diabetes mellitus, but not in adult with insulindependent diabetes. Also diabetes in juvenils is associated with other immune disease like thyroiditis, chronic active hepatitis, infections and thymus deficiency. They suspected that thymus deficiency and autoimmunity play a main role in the pathogenesis of some types of juvenile-onset diabetes mellitus. Moreover, they reported that excess morbidity of the IgAdeficient in juvenile diabetic patients can be due to IgA deficiency in older insulindependent diabetic patients (29). Segade et al, showed patients with type 1 and 2 diabetes have higher level of circulating IgA. Also this phenomenon is more among in male than female (30). Imagava et al, reported that high level of serum enterovirus IgA antibodies in type 1 diabetic patients is due to recurrent enterovirus infection in them (31). Figueredo et al, in one study reported that increased level of immunoglobulin A in diabetic patients suggested a possible role of IgA in the pathogenesis of the vascular complications of diabetes mellitus (32).

Salivary immunoglobulin A and oral complications

It has been shown that salivary component detection in patients with diabetes may help to

References

- Sheikhpour R. Diabetes and Oxidative Stress: The Mechanism and Action. Iran J Diabetes Obes 2013;5(1),40-5
- 2. Sheikhpour R. Glucagon-like peptide 1 and type 2 diabetes: Targets and new therapies. Indian J Sci Technol 2012;5(6):2891-9.
- 3. 3-Sheikhpour R. Evaluation of lipid profile in diabetic patients. Clin Biochem 2011;44(13):160
- Sheikhpour R, Jalali B, Afkhami-Ardekani M. The effect of zinc on lipid oxidizability in diabetic patients in Yazd city. Feyz 2009;13(2):103-10. (in Persian)
- Sheikhpour R, Khoradmehr A. New Insights into the Effect of Diabetes and Obesity in Alzheimer's Disease. Iran J Diabetes Obes 2014;6(1):41-7
- Benfield T, Jensen JS, Nordestgaard BG. Influence of diabetes and hyperglycaemia on infectious disease hospitalisation and outcome. Diabetologia. 2007;50:549-54.
- Leegaard A, Riis A, Kornum JB, Prahl JB, Thomsen VØ, Sørensen HT, et al. Diabetes, glycemic control, and risk of tuberculosis: a population- based case-control study. Diabetes Care. 2011;34:2530-5.
- P. Bakianian Vaziri, M. Vahedi, H. Mortazavi, Sh. Abdollahzadeh, M. Hajilooi. Evaluation of Salivary Glucose, IgA and Flow Rate in Diabetic Patients: A

diagnosis, prevention and management of the manifestations (28).immunoglobulin antibody (S-IgA) plays a main role in the oral homeostasis. The S-IgA is the widespread immunoglobulin in mixed saliva and is assumed to be an important factor for adaptive immunity in the oral cavity. Thus it works as an index of the adaptive local immunity of mouth. S-IgA synergetically with other antibacterial factors, including lysosym, lactopherin, peroxidase, mucine, etc. Thus they prevent the piercing of antigens via the oral mucous membrane (33).

Conclusion

Diabetes mellitus causes an increased risk of morbidity because of infection disease. It seems that low s-IgA levels may be a mechanism to explain susceptibility of infection in DM patients. Early detection of immunoglobulin A deficiency in diabetic patients can reduces risk for morbidity.

- Case-Control Study. J Dentistry, Tehran Univ Med Sci 2010:7(1):13-9
- Oikawa J, Ukawa Sh, Ohira H, Kawamura T, Wakai K. Diabetes Mellitus is Associated With Low Secretion Rates of Immunoglobulin A in Saliva. J Epidemiol 2015;25(7):470-4
- 10. Kakoei Sh, Hosseini B, Haghdoost AA, Sanjari M. Evaluation of Salivary Secretory Immunoglobulin A Levels in Diabetic Patients and Association with Oral and Dental Manifestations. Sultan Qaboos Univ Med J 2015;15(4):507-11.
- Ahmadiafshar A, Mohsenifard M.R, Mazloomzadeh S. Evaluation of Serum & Salivary IgA in Patients with Type 1 Diabetes. Plos one 2015;13:1-7
- 12. Kerr MA. The structure and function of human IgA. Biochem J 1990;271:285-96.
- van Egmond M, van Garderen E, van Spriel AB. FcáRI-positive liver Kupffer cells: reappraisal of the function of immunoglobulin A in immunity. Nat Med 2000:6:680-5.
- 14. Ana Paula França Mantovani1, Mariel Perini Monclaro1, Thelma L Skare. Prevalence of IgA deficiency in adult Systemic Lupus Erythematosus and the study of the association with its clinical and autoantibody profiles. Bras J Rheumatol 2010;50(3):273-82

- Sheikhpour R. Incretins, dipeptidyl peptidase-4 and diabete.ISBN:978-3-659-22207-8. Lambert academic publishing. 2012
- 16. Luciana Salles Branco-de-Almeida, Cláudia Maria Coêlho Alves, Fernanda Ferreira Lopes, Adriana de Fátima Vasconcelos Pereira, Rosane Nassar Meireles Guerra, Antônio Luiz Amaral Pereira. Salivary IgA and periodontal treatment needs in diabetic patients. Braz Oral Res 2011;25(6):550-5
- Bokor-Bratić M, Clinical significance of analysis of immunoglobulin A levels in saliva. Med Pregl 2000;53(3-4):164-8.
- Domenico Greco, Filippo Maggio. Selective Immunoglobulin A Deficiency in Type 1 Diabetes Mellitus: A Prevalence Study in Western Sicily (Italy). Diabetes Metab J 2015;39:132-6
- Salavoura, I Koliaex A, Tsangaris G, Mavvrou A.
 Development of Cancer in Patients with Primary Immunodeficiencies. Anticancer Res 2008;28:1263-70
- 20. Grumach AS, Jacob CMA, Pastorino AC. Deficiência de IgA: avaliação clínico-laboratorial de 60 pacientes
- Latiff A HA, Kerr MA. The clinical significance of immunoglobulin A deficiency. Ann Clin Biochem 2007;44:131-9.
- Wang N, Shen N, Vyse TJ, Anand V, Gunnarson I, Sturfelt G, et al. Selective IgA deficiency in autoimmune diseases. Mol Med 2011;17:1383-96.
- 23. Yel L. Selective IgA deficiency. J Clin Immunol 2010;30:10-6.
- JF, Neovius M, Hammarstrom L. Association between IgA deficiency & other autoimmune conditions: a population-based matched cohort study. J Clin Immunol 2014;34:444-51
- Jonas F. Ludvigsson, Neovius M, Ye W. IgA Deficiency and Risk of Cancer: A Population-Based Matched Cohort Study. J Clin Immunol 2015;35:182-8
- Alkhairy O. IgA Deficiency and Other Immunodeficiencies Causing Mucosal Immunity Dysfunction. Mucosal Immunology 2015;2:1441-9

- 27. Sayarifard F, Aghamohammadi A, Haghi-Ashtiani MT, Rajab A, Irani H, Ahmadian JH, et al. Evaluation of serum IgA levels in Iranian patients with type 1 diabetes mellitus. Acta Diabetol 2012;49(2):131-5
- Mohiti-Ardekani A, Karbassi MH, Mohiti-Ardekani J, Akhondinasab F, Mohammad MH.
 Evaluation of salivary IgA in diabetic and non-diabetic patients: A case-control study. Iran J Diabetes Obes 2012;4:167-71.
- William I Smith Jr, Bruce S Rabin, Alan Huellmantel, David H Van Thiel, Allan Drash. Immunopathology of Juvenile-onset Diabetes Mellitus: I. IgA Deficiency and Juvenile Diabetes. Diabetes 1978;27(11):1092-7
- Segade SR, Camina F, Carnero A, Lorenzo MJ, Alban A, Quinterio C, et al. High serum IgA concentrations in patients with diabetes mellitus: Agewise distribution and relation to chronic complications. Clinical Chemistry 1996;42(7):1064-7
- 31. Imagawa A, Hanafusa T, Makino H, Miyagawa JI, Juto P. High titres of IgA antibodies to enterovirus in fulminant type-1 diabetes. Diabetologia 2005;48:290-3
- 32. Figueredo A, Ibarra JL, Rodriguez A, Molino AM, E.Gomez-de la Concha, Fernandez-Cruz A, et al. Increased Serum Levels of IgA Antibodies to hsp70 Protein in Patients with Diabetes Mellitus: Their Relationship with Vascular Complications. Clinical Immunology and Immunopathology 1996;79(3):252-5
- 33. Rashkova M, Baleva M, Peneva M, Toneva N, Marieta Belcheva M, Koprivarova K. Secretory Immunoglobulin A (S-IgA) and the Oral Risk Markers: Quality of Saliva, Dental Biofilm, Oral Candida and Lactobacillus spp. OHDMBSC 2009;8(3):11-9