Evaluation of CPITN and BMI Indices in Patients with Uncontrolled Type 2 Diabetes

Samira Hajimaghsoodi¹, Akram Ghadiri-Anari², Fahimeh Rashidi-Meybodi³, Zahra Jafarzadeh*⁴

- 1. Assistant Professor, Department of Oral Medicine, School of Dentistry, Shahid Sadoughi University of Medical Sciences, Yazd, Iran.
- 2. Associate Professor, Diabetes Research Center, Shahid Sadoughi University of Medical Sciences, Yazd, Iran.
- 3. Associate Professor, Department of Periodontics, School of Dentistry, Shahid Sadoughi University of Medical Sciences, Yazd, Iran.
- 4. Dental Student, School of Dentistry, Shahid Sadoughi University of Medical Sciences, Yazd, Iran.

*Correspondence:

Jafarzadeh Zahra, Dental Student, School of Dentistry, Shahid Sadoughi University of Medical Sciences, Yazd, Iran.

Tel: (98) 939 443 8960 **Email:** z.smile1992@yahoo.com

Received: 12 October 2017

Accepted: 07 January 2018

Published in February 2018

Abstract

Objective: Diabetes mellitus has multiple effects on oral tissues. Oral symptoms especially periodontal disease and recurrent bacterial and fungal infections seem to vary in patients with different body mass index. The aim of this study was to investigate periodontal treatment needs Index in patients with uncontrolled type 2 diabetes and its relation with body mass index (BMI).

Materials and Methods: In this cross-sectional analytical study, 150 individuals with uncontrolled type 2 diabetes referred to Yazd Diabetes Research Center in 2016 were studied in terms of BMI and the Community Periodontal Index of Treatment Needs (CPITN). Patients divided in two groups: BMI of over 30 and under 30. Data obtained from each group analyzed using SPSS software version 16 and Mann-Whitney statistical test was used.

Results: The mean of CPITN index was not significantly different in men and women. Also in different age groups, there were no significant differences, except 50 to 59 years range. In this age group, the mean of CPITN in patients with BMI above 30 was 1.926 and in BMI under 30 was 1.5792 (*P*-value:0.019). In patients with BMI above 30 the mean of CPITN reported 1.88 in comparison with 1.61 in group with BMI under 30 (*P*-value:0.030).

Conclusion: The results indicated that diabetic patients with a BMI greater than 30 need more periodontal treatment than patient with BMI less than 30. Also indexes in the age range 50 to 59 years was more prominent which means clinicians have to consider patients with uncontrolled type2 diabetes more carefully for providing oral health when BMI and age are increasing.

Keywords: Diabetes, Periodontitis, Body mass index, Community periodontal index of treatment needs

Introduction

iabetes mellitus is the most common endocrine and metabolic diseases. It is determined by an abnormal increase in blood glucose levels and impaired metabolism of carbohydrates, protein and lipid (1). This disease is accompanied by long-term disorders such as retinopathy, nephropathy,

neuropathy, cardiovascular disorders and oral symptoms. The prevalence of diabetes in Iran in 2000 was 7.5% of the population, and by 2025 it will raise to 8.6% that will be equivalent to 1.5 million patient (2-4).

The most important oral symptoms associated with diabetes include dry mouth, burning

mouth, taste changes, candidiasis, increasing the amount of decay and progressive periodontitis. Periodontitis is known as the sixth complication of diabetes and the pathophysiology relationship between them was established. From one side, role of diabetes mellitus as a risk factor for the development of gingivitis and periodontitis is proven and depending on the level of glycemic control, diabetes mellitus increases response to bacterial plaque by gingiva than non-diabetic patients. On the other hand periodontal disease and gingivitis cause weaker control of blood sugar. Also hyperglycemia increase amount of glucose in GCF and causes to change quality of bacteria and increases periodontal diseases in patients with uncontrolled diabetes. (1,5-9)

One of the main risk factors for diabetes mellitus is obesity (2). So far, many studies have investigated a possible link between obesity and periodontal disease. It is thought that hormones and cytokines derived from immune cells in fat tissue cause a series of inflammatory reactions that cause periodontal disease. However, the exact mechanism to determine the relationship between body mass index (BMI) and periodontal disease was not clarified. (10-12)

On the other hand many studies such as Bacic et al. revealed relation between diabetes and loose of teeth which indicates a hypotheses that diabetes can compromise oral health indexes. (13)

In contrast some studies like Rylander et al. with aim to evaluation of periodontal tissues health in diabetes patients and comparison with control group revealed that there were no significant differences between two groups. (14)

Several studies investigated the relationship between the frequency and severity of periodontal disease and obesity or increased BMI in some populations such as heart disease patients, postmenopausal women, people with diabetes and healthy subjects. (12,15-17)

This study designed to evaluate status of periodontal health index in two groups of

uncontrolled type II diabetes regarding to BMI index.

Materials and Methods

This study was an analytical cross-sectional study. According to similar articles the total of 150 samples were used in this study. Patients with uncontrolled type 2 diabetes under treatment with oral hypoglycemic agents referred to Yazd Diabetes Research Center in 2016 which selected by simple random sampling method and divided into two groups. After approval of the project in university ethics committee sampling was done. This research was described initially for patients with written informed consent, and then patients were included in the study. To collect information of patients a checklist consisted of two parts was used, in the first section basic information (including age, sex, and duration of diabetes, the diabetes control level and latest result of HbA1c test) and the other section included BMI and CPITN indexes. BMI indexes were calculated and CPITN indexes were obtained by oral examination with probe and dental mirrors were recorded. Inclusion criteria of patients to the research were uncontrolled type II diabetes and level of latest HbA1c test above or equal to 8, lack of other systemic diseases and age range of 30-59 years old. Exclusion criteria were having less than 20 teeth, currently.

Patients were divided in to two groups based on BMI (BMI < 30, BMI \geq 30). Patients' CPITN index was determined using the 5 codes: (Code 0 = healthy periodontal tissues, Code 1 = bleeding on probing, code 2 = dental calculus, Code 3 = pockets with a depth of less than or equal to 5mm and Code 4 = pocket depth with a depth of 6mm or more) and information was recorded. Sample size was determined according to similar articles considering significant level of 5% and test power of 80%. Data obtained from each group analyzed using SPSS version 16 and Mann-Whitney statistical test was used.

Results

One hundred and fifty samples were divided in two groups of 30≤BMI and 30> BMI. Group A (30≤BMI), including 42 women and 33 men in the age range of 59-30 and an average age of 45 and Group B (30> BMI) included of 23 females and 52 males in the age range of 59-30 and the average age of 40.

The mean of CPITN index in this study did not show significant differences between men and women. (Table 1)

Also for evaluation and comparing of periodontal health and BMI between two groups with 30≤BMI and 30>BMI, group with BMI above 30 and the average of 1.8863 and group with BMI under 30 and the average of 1.6189 revealed significant difference. (Table 2)

The evaluation of periodontal health indexes in different age groups of 30 to 39, 40 to 49

and 50 to 59 revealed that just in group of 50 to 59 years there are significant differences between two group with BMI above 30 and BMI under 30. (Table 3) (*P*-value=0.019)

Also two groups were compared in term of age, duration of diabetes, frequency of teeth brushing and HbA1C test as confounding variables which revealed no significant result for any one. (Table 4)

Discussion

The present study aimed to evaluate and comparison of periodontal status in two groups of patients with uncontrolled diabetes grouping by BMI index.

Prevalence of diabetes in Iran was 5.7% in 2000 and it is estimated to increase up to 6.8% in 2025 which means approximately about 5.1 million people. (3,4)

One of the main risk factors for diabetes

Table 1. Comparing the mean and median CPITN in subjects according to sex

Tubici. Comparing the mean and meanin Cliff, in subjects according to sex							
		CPITN					
Sex	Median	Standard Deviation (SD)	Number	Mean			
Female	1.50	0.75	65	1.57	0.27		
Male	1.50	0.93	85	1.64			
Total	1.50	0.85	150	1.61			

Mann-Whitney Test

Table 2. Comparing of CPITN in two studied groups based on BMI

DMI		CPITN				
BMI	N	Mean	Median	SD		
BMI > 30	75	1.88	1.80	0.671	0.03	
BMI < 30	75	1.61	1.60	0.746		

Mann-Whitney Test

Table 3. Comparing each group in age and CPITN

CPITN									
Age	BMI > 30			BMI < 30				P-Value	
	Mean	Median	SD	N	Mean	Median	SD	N	
30-39	1.61	1.550	0.007	8	1.69	1.800	0.45	6	0.85
40-49	1.90	1.60	0.83	22	1.72	1.800	0.59	16	0.71
50-59	1.92	1.83	0.60	45	1.57	1.400	0.81	53	0.01

Mann-Whitney Test

Table 4. Comparing the mean of age, tooth brushing number, duration of diabetes and HbA1C in two groups

monte in two groups				
Variable	BMI ≥30	BMI< 30	Total	D volue
variable	Mean± SD	Mean±SD Mean ±SD		<i>P</i> -value
Age	50.52 ±7.202	51.73 ±7.282	51.13 ±7.245	0.222
How many toothbrushes a day	2.29 ±1.691	2.27 ± 1.655	2.28 ±1.667	0.952
Duration of disease since diagnosis	9.53 ±6.550	8.68 ± 5.994	9.11 ±6.271	0.466
HbA1c	9.439 ±1.248	9.433 ±1.495	9.436 ±1.372	0.437

Mann-Whitney Test

mellitus is obesity. By raise in body weight occurrence of type II diabetes increases significantly. On the other hand, occurrence of periodontal disease is significantly higher in diabetic patients comparing to non-diabetic patients. (2,18-20)

Distortion and changes in vascularity of gingiva, reduction of immunological response to pathogens in defensive cell lines of host and increase in salivary glucose are assumed as effective factors in occurrence of almost all periodontal diseases in diabetic patients. (15,21-23)

In the present study we aimed to examine the impact of factors such as weight gain on periodontal health in patients with diabetes that may address patients to reach improvements in their periodontal health status. The results of study showed that significant correlation exists between BMI and index of periodontal health in uncontrolled type 2 diabetes patients. In the group with BMI over 30, periodontal health index demonstrated the need to further periodontal treatment.

Many studies have examined the relationship between obesity and periodontal disease.(5,15,19-22,24-27). Most of these studies were performed in healthy subjects in terms of systemic health.(5,15,20-22,25-27)

These results are similar to results of the study of Lalit Kumar Mathur and colleagues which demonstrated that obesity is a risk factor for periodontal disease (20). Studies of Pataro Al et al (5), Andrea Garman et al, and yagob salekzamani et al (22) also confirm the results of the present study.

Vinaly Humar and colleagues in a study in 2013, aimed to evaluate the association between body mass index and periodontal status. it was demonstrated that increase in BMI is a risk factor for periodontitis (25).

In the study by Altamash M and colleagues in 2013, they evaluated the periodontal condition based on HbA1C, BMI and WC (waist circumference) in diabetic patients. Weight gain was observed among all the samples. Pocket depth greater than 6 mm observed in

the sample that had poor glycemic control compared to others and people with poorly controlled diabetes showed severe conditions in periodontal disease (28).

The other study in 2014 aimed to evaluate the relationship between obesity and oral diseases. In that study, oral hygiene index were used which was 13.1 ± 16.2 among overweight patients and $13/1 \pm 5/2$ in normal individuals. It was concluded that there was no significant relationship between increased BMI and periodontal status and BMI only could be a predictor of increase in gingival bleeding (27). In the study about body mass index and periodontitis in patients with type 2 diabetes in Arabs were investigated. 112 samples with 30≤BMI and 74 samples with BMI≤30 were collected. All samples were diabetic and assessed with HbA1C. The study revealed that there was no association between periodontitis and BMI in people with type 2 diabetes, and the authors stated that more studies are needed in this field (24). The study authors noted that the results of the study may be due to sample youth that camouflaged effects of obesity on periodontitis and also due to the index used for identifying periodontitis that pocket depth had been considered, but as we know in some cases presence of periodontitis is without development of pockets.

So far, many hypotheses about the relationship between obesity and periodontal disease have been reported. It is thought that hormones and cytokines derived from immune cells in fat tissue cause a series of inflammatory reactions that cause periodontal disease. However, the exact mechanism to determine the relationship of BMI with periodontal disease has not been revealed. Some probable theories about a possible link between obesity and periodontal disease are: adipose tissue-derived hormones and cytokines, imbalances in the immune system, reduced blood flow in periodontal tissues, hyperlipidemia, and mechanisms related to unhealthy lifestyle and emotional and mental stresses. (29-35)

The results of presented study also showed that only in the age group of 50 to 59 years,

with an increase in BMI, CPITN index also increased. The results are valuable due to its emphasis on the requirement of oral professional care in patients with higher BMI. On the other hand, results in younger patients were not comparable precisely due to the low number of samples, so it might be significant with further studies with higher number of patients.

Also two groups were compared in terms of age, duration of diabetes, frequency of teeth brushing and HbA1C test as confounding variables which revealed no significant result. This showed that there is a normal distribution in samples of the groups which causes higher

research integrity in evaluation of BMI and CPITN indexes.

Conclusions

The results indicate that obese diabetic patients need more periodontal treatment than patients with BMI less than 30. Also indexes and differences in the age range 50 to 59 years is more prominent which means clinicians have to consider patients with uncontrolled type2 diabetes more carefully for providing oral health when BMI and age are increasing.

References

- Burket LW, Greenberg MS, Glick M, Ship JA. Burket's oral medicine: PMPH-USA; 2008.
- Azhdari AE. Check the status of DMFT index in patients with diabetes type II and its relationship with HbA1c in the city of Yazd in 92-1391. Theses.dental College. Yazd University of Medical Sciences martyr. 1392.
- 3. Esteghamati A, Gouya MM, Abbasi M, Delavari A, Alikhani S, Alaedini F, et al. Prevalence of diabetes and impaired fasting glucose in the adult population of Iran. Diabetes care. 2008;31(1):96-8.
- Shaw JE, Sicree RA, Zimmet PZ. Global estimates of the prevalence of diabetes for 2010 and 2030. Diabetes research and clinical practice. 2010;87(1):4-14.
- Pataro AL, Costa FO, Cortelli SC, Cortelli JR, Abreu MHNG, Costa JE. Association between severity of body mass index and periodontal condition in women. Clinical oral investigations. 2012;16(3):727-34.
- 6. Malani PN. Harrison's principles of internal medicine. JAMA. 2012;308(17):1813-4.
- 7. Fauci AS. Harrison's principles of internal medicine: Mcgraw-hill New York; 1998.
- 8. Newman MG, Takei H, Klokkevold PR, Carranza FA. Carranza's clinical periodontology: Elsevier health sciences; 2011.
- Lindhe J, Karring T, Lang NP. Clinical periodontology and implant dentistry: Blackwell Munksgaard Copenhagen; 2003.
- Suvan J, D'Aiuto F, Moles DR, Petrie A, Donos N. Association between overweight/obesity and periodontitis in adults. A systematic review. Obesity Reviews. 2011;12(5):e381-e404.
- 11. Boesing F, Patino J, Da Silva V, Moreira E. The interface between obesity and periodontitis with

- emphasis on oxidative stress and inflammatory response. Obesity reviews. 2009;10(3):290-7.
- 12. Preshaw P, Alba A, Herrera D, Jepsen S, Konstantinidis A, Makrilakis K, et al. Periodontitis and diabetes: a two-way relationship. Diabetologia. 2012;55(1):21-31.
- Bačić M, Plančak D, Granić M. CPITN assessment of periodontal disease in diabetic patients. Journal of Periodontology. 1988;59(12):816-22.
- Rylander H, Ramberg P, Blohme G, Lindhe J. Prevalence of periodontal disease in young diabetics. Journal of clinical periodontology. 1987;14(1):38-43.
- Ekuni D, Mizutani S, Kojima A, Tomofuji T, Irie K, Azuma T, et al. Relationship between increases in BMI and changes in periodontal status: a prospective cohort study. Journal of clinical periodontology. 2014;41(8):772-8.
- 16. Deshpande K, Jain A, Sharma R, Prashar S, Jain R. Diabetes and periodontitis. Journal of Indian Society of Periodontology. 2010;14(4):207.
- 17. Morrison HI, Ellison LF, Taylor GW. Periodontal disease and risk of fatal coronary heart and cerebrovascular diseases. European Journal of Cardiovascular Risk. 1999;6(1):7-11.
- 18. Ghorbani M. A review of the relationship between type 2 diabetes and obesity.
- 19. Al Habashneh R, Azar W, Shaweesh A, Khader Y. The relationship between body mass index and periodontitis among postmenopausal women. Obesity research & clinical practice. 2016;10(1):15-23.
- Mathur LK, Manohar B, Shankarapillai R, Pandya D. Obesity and periodontitis: A clinical study. Journal of Indian Society of Periodontology. 2011;15(3):240.

- Salekzamani Y, Shirmohammadi A, Rahbar M, Shakouri S-k, Nayebi F. Association between human body composition and periodontal disease. ISRN dentistry. 2011;2011.
- 22. Gorman A, Kaye EK, Apovian C, Fung TT, Nunn M, Garcia RI. Overweight and obesity predict time to periodontal disease progression in men. Journal of clinical periodontology. 2012;39(2):107-14.
- 23. Noguchi E, Kato R, Ohno K, Mitsui A, Obama T, Hirano T, et al. The apolipoprotein B concentration in gingival crevicular fluid increases in patients with diabetes mellitus. Clinical biochemistry. 2014;47(1):67-71.
- 24. Awad M, Rahman B, Haidar Hasan HA. The relationship between body mass index and periodontitis in Arab patients with type 2 diabetes mellitus. Oman medical journal. 2015;30(1):36.
- 25. Bhardwaj VK, Sharma D, Jhingta P, Fotedar S, Sahore M, Manchanda K. Assessment of relationship between body mass index and periodontal status among state government employees in Shimla, Himachal Pradesh. Journal of International Society of Preventive & Community Dentistry. 2013;3(2):77.
- Weinspach K, Staufenbiel I, Memenga-Nicksch S, Ernst S, Geurtsen W, Günay H. Level of information about the relationship between diabetes mellitus and periodontitis-results from a nationwide diabetes information program. European journal of medical research. 2013;18(1):6.
- Sede M, Ehizele A. Relationship between obesity and oral diseases. Nigerian journal of clinical practice. 2014;17(6):683-90.
- 28. Altamash M, Arledal S, Klinge B, Engström P-E. Pre-diabetes and diabetes: Medical risk factors and periodontal conditions. Acta Odontologica Scandinavica. 2013;71(6):1625-31.

- 29. Groves DW, Krantz MJ, Hokanson JE, Johnson LR, Eckel RH, Kinney GL, et al. Comparison of frequency and duration of periodontal disease with progression of coronary artery calcium in patients with and without type 1 diabetes mellitus. The American journal of cardiology. 2015;116(6):833-7.
- 30. Matsushita K, Hamaguchi M, Hashimoto M, Yamazaki M, Yamazaki T, Asai K, et al. The novel association between red complex of oral microbe and body mass index in healthy Japanese: a population based cross-sectional study. Journal of clinical biochemistry and nutrition. 2015;57(2):135-9.
- 31. Palle AR, Reddy CSK, Shankar BS, Gelli V, Sudhakar J, Reddy KKM. Association between obesity and chronic periodontitis: a cross-sectional study. The journal of contemporary dental practice. 2013;14(2):168.
- 32. Chaffee BW, Weston SJ. Association between chronic periodontal disease and obesity: a systematic review and meta-analysis. Journal of periodontology. 2010;81(12):1708-24.
- Anand N, Suresh M, Chandrasekaran S. Effect of obesity and lifestyle on the oral health of pre adolescent children. Journal of clinical and diagnostic research: JCDR. 2014;8(2):196.
- Jaramillo A, Lafaurie GI, Millán LV, Ardila CM, Duque A, Novoa C, et al. Association between periodontal disease and plasma levels of cholesterol and triglycerides. Colombia Médica. 2013;44(2):80-6.
- 35. Saito T, Shimazaki Y, Koga T, Tsuzuki M, Ohshima A. Relationship between upper body obesity and periodontitis. Journal of dental research. 2001;80(7):1631-6.