

The Effect of Endurance Training Intensity on the Expression of Perlipin a Protein of Subcutaneous Adipose Tissue and Pancreatic B-cells Function in Diabetic Rats

Hadi Ghaedi*

1. Assistant Professor in Exercise Physiology, Lamerd Branch, Islamic Azad University, Lamerd, Iran.

***Correspondence:**

Hadi Ghaedi, Department of Exercise Physiology, Lamerd Branch, Islamic Azad University, Lamerd, Iran.

Tel: (98) 917 782 8667

Email: ghaedi.hadi@gmail.com

Received: 24 August 2019

Accepted: 05 December 2019

Published in January 2020

Abstract

Objective: The aim of this study was to compare the three endurance training intensities on the expression of Perlipin A protein in subcutaneous adipose tissue and pancreatic B-cells function in male diabetic rats.

Materials and Methods: In this study, 40 healthy male wistar rats were divided into five groups, including diabetic with low intensity endurance training, diabetic with moderate intensity endurance training, diabetic with high intensity endurance training, diabetic control and healthy control. After diabetic induction with streptozotocin, endurance training was performed with low intensity, moderate and severe for eight weeks, three sessions per week. The relative expression of Perilipin A was measured by western blot technique.

Results: The results indicated a significant effect of endurance training with three intensities on serum levels of insulin and glucose and pancreatic B-cells function (P -value: 0.001). Also the results showed that there was no significant difference between Perlipin A expression in healthy and diabetic control groups with endurance training groups (with low, moderate and high intensity) (P -value: 0.07).

Conclusion: However, Moderate and high intensity endurance training compared to low-intensity training can compensate for the loss caused by diabetes in the expression of the Perlipin A protein but the difference was not significant. It seems that more intensity endurance training lead to more increase in Perlipin A expression in diabetic rats.

Keywords: Perlipin A, Endurance training, subcutaneous adipose tissue, Diabetes, pancreatic B-cells function

Introduction

Longitudinal studies have shown that the progression of pancreatic beta cells dysfunction is very important in the

prevalence of diabetes (1,2). Adipose tissue has an important role in controlling whole-body glucose homeostasis in both normal and

disease status (3). SCAT cells may act as a buffer or sink for circulating FFAs and TGs, but once they reach their capacity they lose their protective benefit, fat begins to accumulate in tissues not suited for lipid storage (4). However, most research has pointed to the importance of visceral adipose tissue in glucose metabolism and insulin resistance (5,6,8).

In adipocytes, triglyceride is generally stored in lipid droplets (a neutral lipid core surrounded by a phospholipid monolayer coated with proteins). Positioned at the lipid droplet surface, PAT proteins manage access of other proteins (lipases) to the lipid esters within the lipid droplet core and can interact with cellular machinery important for lipid droplet biogenesis. Genetic variations in the gene for the best-characterized of the mammalian PAT proteins, perilipin, have been associated with metabolic phenotypes, including type 2 diabetes mellitus and obesity (10). Mammalian LDs contain a specific set of proteins called the PAT domain family that include perilipin, adipose differentiation-related protein (ADRP, also called adipophilin (plin2)), a tail-interacting protein of 47 kDa (TIP47) (plin3), S3-12, and OXPAT (plin5) sharing regions of conserved sequences, named PAT domains, mainly at their N termini (11). Perilipin A is one of the most abundant of these proteins, which is activated by protein kinase A, and by controlling various proteins, plays a pivotal role in regulating lipid metabolism in adipocytes (12). Destruction of Perilipin A from white adipose tissue causes adipocyte fat storage disorder by increasing basal lipolysis and reducing the lipolysis stimulated by PKA and will lead to a significant reduction in white adipose tissue (13). The role of Perilipin A in WAT is lipolysis suppression in the absence of PKA stimulation and an increase in lipolysis (about 100 times) with PKA stimulation (14). Data show that modulation of lipid droplet proteins in white adipocytes is a potential therapeutic strategy for the treatment of obesity and

related disorders, such as decreasing blood glucose levels and insulin resistance (15).

Physical activity has a beneficial effect on insulin sensitivity in normal as well as insulin resistant populations. (16). In terms of pancreatic function, experiments in humans and animals have variously demonstrated that exercise improves insulin resistance, increases insulin sensitivity, increases pancreatic beta-cell mass and generally enhances beta cell function and insulin tropic action, especially in type 2 diabetes (17). Sport activity can also increase the mass of the pancreatic beta cells and generally improve and enhance the function of the beta cells of the pancreas (18). Long-term exercise is a common method used to improve health and prevent disease. While physiological compatibility with long-term exercise has been well proven in many devices, such as skeletal muscle, certain changes that occur under adipose tissue, especially subcutaneous adipose tissue, are underestimated (19). Its exact mechanism for lipolysis regulation is not completely determined (20). Many sources report enhanced stimulated lipolysis in adipocytes from endurance trained subjects that is attributed to alterations in the impolitic cascade. The purpose of this study was to compare the severity of endurance training on the expression of perilipin A protein of subcutaneous adipose tissue and pancreatic B-cells function in STZ-induced diabetic rats.

Materials and Methods

Animals

In this study, 48 eight-week-old healthy male wistar rats (250 ± 40 gr) were used in the present study purchased from Pasture Institute (Tehran, Iran). The animals were housed two per polypropylene cages at a room temperature of $22\text{ }^{\circ}\text{C}$ with 12:12h dark: light cycle (lightings were turned on at 8 PM and off at 8 AM) and 60% air humidity with free access to water and rodent pellet food. Rats were divided into the following five groups randomly: the low intensity training group (n= 8), moderate intensity training group (n= 8),

high intensity training group (n= 8), diabetic control group (n= 8) and healthy nondiabetic sedentary group (n= 8).

Diabetes was induced by intraperitoneal injection of a single dose of streptozotocin in a fasting condition (STZ, 55 mg/ body weight; Sigma, St Louis, MO, USA) (28). To reduce the effect of streptozotocin on beta cells 15 minutes before streptozotocin injection, 270 mg / kg of nicotinamide saline mice was used (29). Rats in sham just received buffer. Two days after STZ injection blood was obtained from a small nick in the tail and measured with a glucometer (Optium Xceed). Rats with a glucose concentration exceeding 300 mg/dL were considered as a hyperglycemic state (30). From the diabetic experimental group comprising 40 rats, four rats in whom diabetes was not induced and six rats in poor health were excluded.

Wistar rats carried out eight-week treadmill training with different intensities for 30 min/training and 4 days per week for 8 weeks on a standard treadmill for small rodents. Rats were divided into groups including low intensity training group (5-8 m/min, 50-60% VO₂max), moderate exercise intensity group (14-16m/min, 65-70% VO₂max), and high intensity exercise group (22-25 m/min, 80% VO₂max) (31). In addition, voluntary continuous exercise was induced by a 10-voltage electronic stimulator. After one week and familiarity with the laboratory environment, the rats were first introduced to the rats by running on a treadmill for a week at a speed of 10 m / min for fifteen to twenty minutes of training (32).

At the end of eight weeks of endurance training in order to rule out temporary training effects of last session, tissue sampling was conducted 48 h after last exercise bout. All rats were fasted overnight by removing food from the rat's cages 12 h prior to sacrifice. They were sacrificed for tissue collection by an intraperitoneal injection of ketamine (75 mg/kg) and xylazine (25 mg/kg). The adipose tissue was excised, frozen in liquid nitrogen and stored frozen at -80°C until further

analysis. The serum glucose levels were measured by glucometer in Germany, by measuring the tip of the tail and serum levels of insulin with a rat specific ELISA kit (Insulin rat ELISA DEV8811) manufactured by Demeditec in Germany with a sensitivity of 0.1 ng / mL and pancreatic B-cells function using the following formula was obtained (33).

FPI=Fasting Plasma Insulin (ng/ml)

FPG=Fasting Plasma Glucose (mg/dl)

HOMA-B= (20×FPI)/ (FPG- 3/5)

In order to analyses of visceral adipose tissue protein expression of Prilipin A by western blot, approximately 50 mg of each visceral adipose tissue pieces was powdered with a pestle in liquid nitrogen and lysed using a 1 mL of phosphate-buffered saline (PBS).

Statistical analysis

Data are reported as means ±standard error (SE) values. To compare groups, dependent variables were analyzed by One-way variance analysis and Kruskal–Wallis. The post hoc Games-Howell test was used to determine significant differences between groups. Statistical significance was set at P -value< 0.05.

Ethical considerations

This research was experimental. Animal procedures were approved by Shahrekord university ethics committee of animals (SKU: MP221/2013).

Results

The results of one-way ANOVA showed significant effect of exercise on serum levels of insulin and glucose and pancreatic B-cells function (P -value= 0.001) (Table 1), However, the results of the post hoc Games-Howell test showed that pancreatic B-cells function was significant different between experimental groups and healthy control group (P -value: 0.001).

Comparison of protein expression of perilipin A by using Kruskal-Wallis test in five groups showed no significant difference (P -value= 0.07).

Table 1. Compare Serum concentration of glucose and insulin, pancreatic B-cells function and perilipin A among groups

Variable	Groups	HC	DC	DHE	DME	DLE
Glucose (mg/ dl)	Mean (\pm SD)	156.25 (\pm 22.94)	557.75 (\pm 158.84)	341.500 (\pm 91.90)	323.13 (\pm 48.61)	497 (\pm 25.43)
	F	18.89				
	P-value	*0.001				
Insulin (ng / ml)	Mean (\pm SD)	0.49 (\pm 0.031)	0.19 (\pm 0.03)	0.12 (\pm 0.009)	0.14 (\pm 0.03)	0.17 (\pm 0.03)
	F	211.35				
	P-value	*0.001				
pancreatic B-cells function	Mean (\pm SD)	0.65 (\pm 0.01)	0.007 (\pm 0.003)	0.007 (\pm 0.002)	0.013 (\pm 0.011)	0.007 (\pm 0.001)
	F	98.74				
	P-value	*0.001				
Perilipin A (Relative density of the band)	Mean (\pm SD)	25600 (\pm 11300.08)	7364.50 (\pm 4436.32)	14483.75 (\pm 8539.62)	10857.87 (\pm 5858.42)	8592.45 (\pm 5401.68)
	Chi-Square	8.49				
	Df	1.40				
	P-value	0.07				

HC: Healthy control group[‡] DC: diabetic control group[‡] DLE: diabetic and low intensity training group[‡] DME: diabetic and moderate intensity training group[‡] DHE: diabetic and high intensity training group

The results showed that perilipin A content in training groups was more likely to increase with increasing exercise intensity than diabetic control group however, this increase was not significant. In addition, the density of bands in different groups is shown in Figure 1.

Discussion

Changes in the expression of lipid droplet adipocyte proteins, such as perilipin A cause alter lipolysis and insulin resistance (24). The

findings showed that insulin and glucose levels decreased with increasing intensity of endurance training in diabetic rats. There was no difference between the experimental groups (low, moderate and high intensity exercise) of pancreatic beta cells function and the only difference was observed between the experimental groups and the healthy control group. Reducing serum levels of insulin and glucose in diabetic groups with high intensity exercise and moderate intensity exercise




Figure 1. Changes in expression of perilipin A in groups

compared to diabetic control and low intensity exercise was significant. Possible reducing serum insulin and glucose mechanisms through endurance exercise can include increase carrier proteins of glucose (GLUT4), decrease secretion and increase cleaning free fatty acids, increase glucose delivery to the muscles tendency to available glucose. High intensity and long duration exercise increase insulin sensitivity possibly by increasing the transfer of glucose to the muscle or reducing fatty acids synthesis and glucose reabsorption due to skeletal muscles activity (25). Also decrease in fat mass can be another reason of reducing insulin (26). Most studies reported this decrease index following exercise schedule, have had a relatively high intensity exercise (27,28). Stephan et al. (2017) in a research showed the mean disposition index as a validated measure of β -cell function, was significantly increased after the intervention. Insulin processing inefficiency in the β -cell, expressed as the fasting proinsulin-to insulin ratio, was also reduced and increased β -cell function during the early-phase response to glucose correlated significantly with reductions in abdominal body fat (29). A recently published study of 8 weeks of high-intensity interval training, which has risen in popularity in both the exercise science field and society at large, in adults with type 2 diabetes mellitus showed improvements in β -cell function, despite no changes in insulin secretion or sensitivity (30). While Huang et al. (2011) showed that 6 weeks of voluntary exercise in type 1 diabetic rat increased the content and secretion of insulin, but did not change in the density islets of langerhans and the number of beta cells (31).

The results of some of these studies are consistent with our findings. Regarding these, the decrease in insulin and glucose levels seems reasonable in the present study. Contradiction in studies results can be caused by different factors such as nutrition type, exercise program, subject's type and intensity and duration of exercise.

Also, although there was no significant difference in the expression of perilipin A protein among diabetic training groups (low, moderate, and high). However, with increasing exercise intensity, there was a tendency to increase in perilipin A. It seems that the non-significant increase of perilipin A in diabetic rats has somewhat compensated for the reduction of perilipinA in diabetic control rat and prevents the deleterious effects of increased lipolysis and the release of fatty acids (32). Because inefficiency in lipolytic suppression and Non-esterified fatty acids (NEFA) increase when FFA demand is low, it can have serious metabolic consequences and is believed to be a key mechanism for the development of type 2 diabetes (32). Increasing the nonsignificant expression of the perilipin A in diabetic rats can also be related to the reduction of TNF α (26) and leptin (33), as well as the increase of plasma lactate (34) that may have been induced by exercise. It has been shown that TNF α reduces the expression of perilipin and increases basal lipolysis in vitro. However, the effects of TNF α on perilipinA expression should also be investigated in vivo (35). Ju Yong Bae et al in 2017 showed that after eight weeks of training, the levels of PKA, Perlipin, CGI-58, ATGL and HSL increased (36). Also, Ramos et al. (2016) showed that endurance training did not change the content of PLIN protein in epididymal white adipose tissue (37). The results of this research are consistent with the current research. Observed differences in some of this studies may be relevant to difference of type of tissue (adipose or muscle), type of fat stores (visceral or subcutaneous), type of exercise (acute or chronic), exercise protocol (duration and intensity), subjects (human and animal), type of subjects (healthy or patient).

Conclusions

Endurance training with different intensities led to changes in serum levels of insulin and glucose compared to diabetic control group. But endurance training did not significantly change the pancreatic beta cells function

compared to diabetic control group. However, the difference was significant in comparison with the healthy control group. Also, there was no significant difference in the expression of perilipin A in the groups, but, it seems increasing intensity of endurance exercise training let to more increase in perilipin A expression in diabetic rats. It is likely that the exercise with the appropriate intensity and duration can increase the level of perilipin A,

which can increase insulin sensitivity and improve insulin resistance in diabetic rats.

Acknowledgments

This article is based on a research project with code 2409511120016 was sponsored by Lamerd Islamic Azad University.

Funding

Funding for this research was provided by the Islamic Azad University of Lamerd Branch.

References

1. Weyer C, Bogardus C, Mott DM, Pratley RE. The natural history of insulin secretory dysfunction and insulin resistance in the pathogenesis of type 2 diabetes mellitus. *The Journal of clinical investigation*. 1999;104(6):787-94.
2. Powelka AM, Seth A, Virbasius JV, Kiskinis E, Nicoloso SM, Guilherme A, et al. Suppression of oxidative metabolism and mitochondrial biogenesis by the transcriptional corepressor RIP140 in mouse adipocytes. *The Journal of clinical investigation*. 2006;116(1):125.
3. Guilherme A, Virbasius JV, Puri V, Czech MP. Adipocyte dysfunctions linking obesity to insulin resistance and type 2 diabetes. *Nature reviews Molecular cell biology*. 2008;9(5):367-77.
4. Ibrahim MM. Subcutaneous and visceral adipose tissue: structural and functional differences. *Obesity reviews*. 2010;11(1):11-8.
5. Colberg SR, Simoneau J, Thaete FL, Kelley DE. Skeletal muscle utilization of free fatty acids in women with visceral obesity. *Journal of Clinical Investigation*. 1995;95(4):1846.
6. Ross R, Fortier L, Hudson R. Separate associations between visceral and subcutaneous adipose tissue distribution, insulin and glucose levels in obese women. *Diabetes Care*. 1996;19(12):1404-11.
7. Bickel PE, Tansey JT, Welte MA. PAT proteins, an ancient family of lipid droplet proteins that regulate cellular lipid stores. *Biochimica et Biophysica Acta (BBA)-Molecular and Cell Biology of Lipids*. 2009;1791(6):419-40.
8. Yamaguchi T, Matsushita S, Motojima K, Hirose F, Osumi T. MLDP, a novel PAT family protein localized to lipid droplets and enriched in the heart, is regulated by peroxisome proliferator-activated receptor α . *Journal of Biological Chemistry*. 2006;281(20):14232-40.
9. Braseamble D. The perilipin family of structural lipid droplet proteins: stabilization of lipid droplets and control of lipolysis. *The Journal of Lipid Research*. 2007;48:2547-59.
10. Martinez-Botas J, Anderson JB, Tessier D, Lapillonne A, Chang BH-J, Quast MJ, et al. Absence of perilipin results in leanness and reverses obesity in *Lepr^{db/db}* mice. *Nature genetics*. 2000;26(4):474.
11. Miyoshi H, Souza SC, Zhang H-H, Strissel KJ, Christoffolete MA, Kovsan J, et al. Perilipin promotes hormone-sensitive lipase-mediated adipocyte lipolysis via phosphorylation-dependent and-independent mechanisms. *Journal of Biological Chemistry*. 2006;281(23):15837-44.
12. Sawada T, Miyoshi H, Shimada K, Suzuki A, Okamatsu-Ogura Y, Perfield II JW, et al. Perilipin overexpression in white adipose tissue induces a brown fat-like phenotype. *PLoS one*. 2010;5(11):e14006.
13. Borghouts L, Keizer H. Exercise and insulin sensitivity: a review. *International journal of sports medicine*. 2000;21(01):1-12.
14. Howarth FC, Marzouqi F, Al Saeedi A, Hameed RS, Adeghate E. The effect of a heavy exercise program on the distribution of pancreatic hormones in the streptozotocin-induced diabetic rat. *JOP Journal of the Pancreas (Online)*. 2009;10(5):485-91.
15. Snowling NJ, Hopkins WG. Effects of different modes of exercise training on glucose control and risk factors for complications in type 2 diabetic patients. *Diabetes care*. 2006;29(11):2518-27.
16. Pistor KE. *The Effects of Endurance Exercise Training on Adipose Tissue Metabolism*: York University; 2012.
17. Ceddia RB, Anthony NM, Patel P, Noor F, Hawke TJ, Gaidhu MP. dysregulation of lipolysis and lipid metabolism in visceral and subcutaneous adipocytes by high-fat diet. *The FASEB Journal*. 2010;24(1):892.4-4.
18. Szkudelski T. The mechanism of alloxan and streptozotocin action in B cells of the rat pancreas. *Physiological research*. 2001;50(6):537-46.
19. Novelli M, Poci A, Lajoix A, Befly P, Bezzi D, Marchetti P, et al. Alteration of β -cell constitutive

- NO synthase activity is involved in the abnormal insulin response to arginine in a new rat model of type 2 diabetes. *Molecular and cellular endocrinology*. 2004;219(1):77-82.
20. Pushparaj P, Low H, Manikandan J, Tan B, Tan C. Anti-diabetic effects of *Cichorium intybus* in streptozotocin-induced diabetic rats. *Journal of Ethnopharmacology*. 2007;111(2):430-4.
 21. Kim D-H, Kim S-H, Kim W-H, Moon C-R. The effects of treadmill exercise on expression of UCP-2 of brown adipose tissue and TNF- α of soleus muscle in obese Zucker rats. *Journal of Exercise Nutrition & Biochemistry*. 2013;17(4):199.
 22. Wisløff U, Helgerud J, Kemi OJ, Ellingsen Ø. Intensity-controlled treadmill running in rats: $\dot{V}O_2$ max and cardiac hypertrophy. *American Journal of Physiology-Heart and Circulatory Physiology*. 2001;280(3):H1301-H10.
 23. Wallace TM, Levy JC, Matthews DR. Use and abuse of HOMA modeling. *Diabetes care*. 2004;27(6):1487-95.
 24. Greenberg AS, Coleman RA, Kraemer FB, McManaman JL, Obin MS, Puri V, et al. The role of lipid droplets in metabolic disease in rodents and humans. *The Journal of clinical investigation*. 2011;121(6):2102.
 25. Kim ES, Im JA, Kim KC, Park JH, Suh SH, Kang ES, et al. Improved insulin sensitivity and adiponectin level after exercise training in obese Korean youth. *Obesity*. 2007;15(12):3023-30.
 26. Paman W, Westerterp-Plantenga M, Saris W. The effect of exercise training on leptin levels in obese males. *American Journal of Physiology-Endocrinology And Metabolism*. 1998;274(2):E280-E6.
 27. Ivy JL. Role of exercise training in the prevention and treatment of insulin resistance and non-insulin-dependent diabetes mellitus. *Sports medicine*. 1997;24(5):321-36.
 28. Kelley DE, Goodpaster BH. Effects of physical activity on insulin action and glucose tolerance in obesity. 1999.
 29. Nieuwoudt S, Fealy CE, Foucher JA, Scelsi AR, Malin SK, Pagadala M, Rocco M, Burguera B, Kirwan JP. Functional high-intensity training improves pancreatic β -cell function in adults with type 2 diabetes. *American Journal of Physiology-Endocrinology and Metabolism*. 2017;313(3):E314-20.
 30. Madsen SM, Thorup AC, Overgaard K, Jeppesen PB. High intensity interval training improves glycaemic control and pancreatic β cell function of type 2 diabetes patients. *PloS one*. 2015;10(8).
 31. Huang H-H, Farmer K, Windscheffel J, Yost K, Power M, Wright DE, et al. Exercise increases insulin content and basal secretion in pancreatic islets in type 1 diabetic mice. *Experimental diabetes research*. 2011;2011.
 32. Nielsen TS, Jessen N, Jørgensen JOL, Møller N, Lund S. Dissecting adipose tissue lipolysis: molecular regulation and implications for metabolic disease. *Journal of molecular endocrinology*. 2014;52(3):R199-R222.
 33. Chapados N, Collin P, Imbeault P, Corriveau P, Lavoie J-M. Exercise training decreases in vitro stimulated lipolysis in a visceral (mesenteric) but not in the retroperitoneal fat depot of high-fat-fed rats. *British journal of nutrition*. 2008;100(3):518-25.
 34. Hashimoto T, Sato K, Iemitsu M. Exercise-inducible factors to activate lipolysis in adipocytes. *Journal of Applied Physiology*. 2013;115(2):260-7.
 35. Johnson EA. Regulation of Lipolysis By Perilipin: influence of Obesity and Exercise Training. 2010.
 36. Bae JY, Woo J, Roh HT, Lee YH, Ko K, Kang S, et al. The effects of detraining and training on adipose tissue lipid droplet in obese mice after chronic high-fat diet. *Lipids in health and disease*. 2017;16(1):13.
 37. Ramos SV, Turnbull PC, MacPherson RE. Adipose tissue depot specific differences of PLIN protein content in endurance trained rats. *Adipocyte*. 2016;5(2):212-23.