

Innovations in Functional Food Development for Diabetes Management: A Narrative Review

Fatemeh Ghannadiasl*

Department of Food Sciences and Technology, Faculty of Agriculture and Natural Resources, University of Mohaghegh Ardabili, Ardabil, Iran.

Abstract

This review explores advancements in the food industry for developing functional foods to manage diabetes. It evaluates the mechanisms and clinical evidence for eight bioactive components. Dietary fibers help control blood sugar levels and insulin sensitivity by slowing glucose absorption and promoting the production of short-chain fatty acids. Polyphenols reduce glucose spikes by inhibiting carbohydrate-digesting enzymes and activating signaling pathways like AMPK. Antioxidants protect pancreatic β -cells and insulin secretion by combating oxidative stress and inflammation. Probiotics and prebiotics improve metabolic balance by modulating gut microbiota and strengthening the intestinal barrier. Omega-3 and monounsaturated fatty acids reduce inflammation and enhance insulin receptor signaling. Fermented foods provide bioactive compounds with hypoglycemic effects, while alkaloids, flavonoids, and terpenoids have diverse anti-diabetic properties. Clinical trials have shown promising results in lowering fasting glucose, HbA1c and postprandial spikes, but limitations like small sample sizes and inconsistent formulations call for further validation through robust randomized controlled trials. Conflicting results across studies emphasize the need for cautious interpretation.


Keywords: Diabetes mellitus, Blood glucose, Functional foods

QR Code:



Citation: Ghannadiasl F. Innovations in Functional Food Development for Diabetes Management: A Narrative Review. IJDO 2025; 17 (4) :272-282

URL: <http://ijdo.ssu.ac.ir/article-1-995-en.html>

 10.18502/ijdo.v17i4.20042

Article info:

Received: 8 August 2025

Accepted: 17 September 2025

Published in October 2025



This is an open access article under the (CC BY 4.0)

Corresponding Author:

Fatemeh Ghannadiasl, Department of Food Sciences and Technology, Faculty of Agriculture and Natural Resources, University of Mohaghegh Ardabili, Ardabil, Iran.

Email: ghannadiasl@uma.ac.ir

Orcid ID: 0000-0001-8257-8559

Tell: (98) 453 150 5160

Introduction

Diabetes mellitus is a major worldwide health issue, with its rising rapidly in different parts of the world. It is defined by high blood sugar levels due to the body's inability to produce or use insulin effectively (1). This condition affects approximately 537 million people worldwide and is a leading cause of morbidity and mortality (2). The rising prevalence of diabetes, particularly in low- and middle-income countries, imposes a considerable economic burden on individuals and healthcare systems. This is due to the chronic nature of the disease and its severe macrovascular and microvascular complications (3).

The challenges associated with diabetes management are multifaceted, encompassing socio-cultural, economic, and healthcare system-related barriers. Individuals diagnosed with diabetes often face difficulties adhering to complex self-management practices, which include dietary modifications, physical activity, medication adherence, and regular monitoring of blood glucose levels (4). Epidemiological evidence links elevated blood levels of pro-inflammatory markers to diets high in processed foods, full-fat dairy, refined grains, and sugar-sweetened beverages. Moreover, inflammation correlates not only with these overall eating patterns but also with the specific balance and intake levels of macronutrients and micronutrients (5). Functional foods are rich in bioactive compounds such as polyphenols, flavonoids, and fibers, which have demonstrated hypoglycemic properties (6,7). These foods influence glucose metabolism through multiple mechanisms, including enhancing gut health, modulating the gut microbiota, and improving gut barrier integrity. These effects can lead to improved insulin sensitivity and reduced inflammation, both of which are essential for effective diabetes management (8, 9).

Recent studies highlight the importance of incorporating innovative food technologies to improve glycemic control in individuals with diabetes. Kaur et al. (2021) emphasize that the development of low glycemic index foods and novel functional ingredients can significantly reduce postprandial blood sugar spikes (10). A systematic review and controlled clinical trial demonstrated that the use of soluble fibers (11,12) and low- or non-nutritive sweeteners (13) effectively lowers postprandial glucose and insulin responses, providing a complementary approach to traditional dietary management. Additionally, a systematic review and meta-analysis of randomized controlled trials showed improvements in markers of fasting and postprandial glycemic control with beta-glucan-rich foods (14). In a controlled trial, starch-restricted, fiber-rich functional bread with an increased beta-glucan-to-starch ratio improved long-term metabolic control in diabetic patients (15). Furthermore, a systematic review and meta-analysis of randomized controlled trials suggested that probiotics and synbiotics may be effective in managing diabetes (16). It appears that innovations in functional food development offer a transformative opportunity to optimize glycemic control and improve metabolic outcomes.

Definition of functional foods

Functional foods are defined as novel or conventional foods that provide health benefits beyond basic nutrition when consumed regularly at effective levels (17). These foods include whole foods, as well as fortified, enriched, or enhanced foods that may offer beneficial effects on health. However, the definition of the term remains complex, and scientific investigation is essential to prove the bioavailability and efficacy of these compounds at physiologically achievable levels within typical dietary patterns (18).

It is important to note that functional foods, dietary supplements, and nutraceuticals are

distinct categories situated at the intersection of nutrition and pharmaceuticals, each regulated by unique frameworks (19). Functional foods are regular foods that provide specific health benefits beyond basic nutrition when consumed as part of a regular diet (20). Dietary supplements, including vitamins, and botanical extracts, are used to improve or maintain health (21). Nutraceuticals encompass both functional foods and dietary supplements, offering health benefits that include prevention and treatment. (20). Developing a framework that categorizes functional foods based on volume and quality of scholarly research could enhance public education, inform clinical guidance, and establish functional foods as a reliable strategy for promoting health and well-being (22).

Innovations in the food industry for functional foods

The food industry is increasingly focusing on functional foods, which offer health benefits beyond basic nutrition (23). This trend is driven by growing consumer awareness of the link between diet and health, as well as the potential of food industry by-products as valuable ingredients (24). These by-products are nutrient-rich, contain bioactive compounds, and have functional properties, making them ideal for creating innovative food products (25). However, to ensure a product qualifies as functional, companies must adhere to specific guidelines for development, efficacy testing, and safety assessment (26).

Recent advancements in technologies and production methods have significantly accelerated the development of functional foods for managing diabetes. Fermented products, a prominent class of functional foods, have shown antidiabetic effects in both *in vitro* and *in vivo* studies (27). Similarly, food-derived bioactive peptides produced through enzymatic hydrolysis (top-down) or assembled via synthetic routes (bottom-up) demonstrate promising glycemic regulatory properties. But their clinical benefits require

confirmation through well-powered, large-scale trials (28). Biotechnology has been instrumental in incorporating targeted bioactive compounds and the enhancement of fermentation processes to improve the health-promoting qualities of functional food products (29). Despite these advancements, challenges such as high production costs, limited market access (30), the need for rigorous scientific validation of health claims, and the requirement to offer a variety of products for regular consumption (31) remain unresolved. Key formulation strategies include incorporating of bioactive ingredients, removing undesirable components, adjusting existing constituents to meet market demands, and improving metabolic outcomes in diabetic patients (32).

Bioactive components and their role in diabetes management

Dietary Fibers

Dietary fiber is essential for managing diabetes, due to its bioactive components that impact metabolism. Soluble fiber, found in foods like oats, beans, and fruits, helps control blood sugar levels by slowing glucose absorption and enhancing insulin sensitivity. Insoluble fiber, present in whole grains and vegetables, promotes digestive health and can aid in weight management, which is crucial for diabetes control (33). Additionally, short-chain fatty acids (SCFAs), produced from fiber fermentation in the gut, can improve insulin secretion and β -cell function, benefiting those with type 2 diabetes (34). Phytochemicals and antioxidants in fiber-rich foods may also reduce inflammation and oxidative stress, supporting overall metabolic health (35). Research has shown that increasing dietary fiber intake can lead to significant reduction in HbA1c levels, fasting plasma glucose, and body weight (36).

Polyphenols

Polyphenols, a diverse group of bioactive compounds found in plant-based foods, have attracted significant attention for their

potential role in managing of diabetes. These compounds enhance insulin sensitivity and glucose metabolism, both are crucial for effective diabetes management (37,38). Polyphenols decrease glucose absorption in the intestine and inhibit carbohydrate digestion, resulting in lower blood sugar levels (38,39). They also regulate intracellular signaling pathways, such as activating of AMP-activated protein kinase (AMPK), which plays a role in glucose transport and metabolism (39). Polyphenols complement conventional diabetes treatments, potentially reducing the side effects of pharmaceutical interventions (40). Despite, promising mechanisms like reducing oxidative stress, suppressing pro-inflammatory cytokines, and modulating key metabolic regulators like AMPK and peroxisome proliferator-activated receptors (PPARs) (41), findings from epidemiological and intervention studies are inconsistent (42). Variability in bioavailability and individual metabolic responses (41, 43, 44) likely contribute to this heterogeneity. Therefore, well-designed, standardized intervention trials are necessary to clarify conflicting findings and confirm the anti-diabetic effects of dietary polyphenols.

Antioxidants

The importance of antioxidants and bioactive compounds in managing diabetes is increasingly acknowledged for their ability to combat oxidative stress, a key factor in the development of the disease. These compounds can improve insulin secretion, protect β -cells, and reduce diabetes-related complications. Oxidative stress occur when there is an imbalance between reactive oxygen species (ROS) and antioxidants, leading to complications (45). Antioxidants work by scavenging free radicals, thereby reducing oxidative damage and enhancing metabolic function (46). Flavonoids and carotenoids, known for their antioxidant and anti-inflammatory properties, can boost β -cell function and insulin secretion. They may also prevent β -cell apoptosis and promote

proliferation, offering protection effects against diabetes (47).

While some studies suggest that antioxidant supplements can benefit diabetes complications, other do not show a clear clinical advantage, resulting in conflicting evidence (48,49). A meta-analysis indicated that antioxidant treatment led to significant improvements in fasting glucose, HbA1c and oxidative stress biomarkers (49). However, other reviews highlight inconsistent findings among trials and question the ability of antioxidants to prevent atherosclerosis or reduce clinical complications in individuals with diabetes (50). The effectiveness of antioxidants likely varies based on the specific compound tested, dosage and formulation, study duration, patient characteristics, baseline antioxidant levels, and concurrent treatments (51). Therefore, well-designed randomized trials with sufficient power standardized interventions and relevant endpoints are needed to determine which antioxidants, at what doses and in which patient populations can effectively reduce in diabetes related complications.

Probiotics

Probiotics are being explored as potential aids in managing diabetes management, primarily because they can help regulate gut microbiota and improve metabolic health. Probiotics play a role in restoring the balance of gut microbiota, which is often disrupted in individuals with diabetes, leading to better metabolic outcomes (52,53). Certain probiotics, like *Clostridium butyricum*, produce butyrate, which helps regulate blood sugar levels and reduces inflammation (54). Probiotics also strengthen the intestinal barrier, reducing the risk of metabolic endotoxemia, which can contribute to insulin resistance (55). *Lactobacillus sporogenes* has been linked to improved glycemic responses and overall metabolic health. *Streptococcus faecalis* has been shown to be effective in lowering postprandial glucose levels, particularly in experimental models. *Bacillus*

mesentericus is known for its role in gut health and may contribute to better diabetes management through its probiotic effects (54). However, the evidence supporting the efficacy of probiotics varies significantly depending on the specific strains (56,57), doses, and treatment durations (58). Some show benefits for blood sugar control, while others have minimal effect. Meta-analyses indicate moderate variability in outcomes, suggesting that strain-specific effects play a significant role in the results. This variability underscores the importance of considering strain-specific effects and individual factors before drawing broad conclusions (56).

Prebiotic

Prebiotics, non-digestible carbohydrates, support the growth of beneficial gut bacteria, enhancing insulin sensitivity and glucose metabolism. They influence gut microbiota by promoting the growth of beneficial bacteria, such as butyrate producers, linked to improved glucose metabolism (59). Prebiotic consumption leads to the production of SCFAs, which help lower blood sugar levels and reduce inflammation (60). These effects can enhance insulin sensitivity and reduce insulin resistance, which is crucial in managing diabetes. Inulin, fructooligosaccharides, and galactooligosaccharides, are well-studied prebiotics with significant health benefits (61-62). Tailoring prebiotic intake to individual gut microbiota profiles may further optimize diabetes management strategies (59).

Beneficial fatty acids

Fatty acids are essential in the development of type 2 diabetes as they impact insulin receptor signaling and glucose transporter translocation. The specific composition of dietary fatty acids can worsen or improve insulin resistance, highlighting their significance in diabetes management. It is crucial to be aware that saturated and trans fats can increase insulin resistance and inflammation, which are detrimental to diabetes control (63). Saturated fatty acids

(SFAs) are known to contribute to insulin resistance, a hallmark of type 2 diabetes mellitus. They can disrupt insulin signaling pathways and impair glucose transporter translocation (64). High consumption of SFAs is associated with higher levels of pro-inflammatory cytokines, exacerbating insulin resistance and β -cell dysfunction (63). Conversely, certain dietary fats, such as omega-3 fatty acids, are linked to a lower risk of type 2 diabetes. Research indicates that regular intake of these fats can reduce inflammation and improve insulin sensitivity, thereby reducing the risk of diabetes (65). Current guidelines recommend reducing SFA intake while increasing monounsaturated and polyunsaturated fatty acids consumption to lower the risk of diabetes and enhance the management of existing conditions (66).

Fermented foods

Fermented foods are known to boost antioxidant activity, which can help reduce oxidative stress linked to diabetes (67). Bioactive peptides created during fermentation have antidiabetic properties that can enhance insulin sensitivity and glycemic control (68,69). Additionally, organic acids, like lactic acid, can lower blood sugar levels and promote gut health, which is important for managing diabetes (70). Beneficial bacteria, such as *Lactobacillus* and *Bifidobacterium* species found in fermented foods, can influence the gut microbiota, improve metabolic functions, and potentially assist in weight management (68,71). In a randomized controlled trial, individuals with type 2 diabetes who consumed probiotic fermented milk (kefir) for 8 weeks experienced a notable decrease in fasting blood glucose and HbA1c levels compared to a control group (72). However, the overall health benefits of fermented foods are not yet fully backed by strong clinical evidence, and the effects can vary depending on the specific food, microbial strains, and preparation method (73).

Functional Plant Compounds

These natural substances, derived from various medicinal plants, have different mechanisms that improve insulin sensitivity, decrease glucose absorption and reduce oxidative stress. Key bioactive compounds such as alkaloids, flavonoids, terpenoids, and polyphenols, have shown significant antidiabetic properties (74). They enhance insulin sensitivity, inhibit carbohydrate-digesting enzymes, and regulate glucose metabolism (74,75). While preclinical studies show promise for alkaloids and flavonoids, further clinical trials are needed to determine optimal dosages and potential interactions with other medications (76). Salleh (2021) points out the challenges of human clinical trials involving medicinal plants, and stresses the importance of standardized methodologies and reliable data collection (77). It is crucial to conduct rigorous clinical trials to confirm the efficacy and safety of these compounds (78). Caution is advised regarding potential interactions between flavonoid-containing supplements and medications (79). Therefore, the therapeutic potential of these plant-derived compounds should be interpreted cautiously until supported by high-quality clinical trials.

Mechanisms of action in diabetes management

Reduce glucose absorption and improve insulin function

The impact of functional foods on blood glucose levels is significant, especially in reducing glucose absorption and improving insulin activity. Functional foods that are high in polysaccharides and dietary fiber can slow down glucose absorption in the intestines, resulting in lower postprandial blood sugar levels (7). Foods with a low glycemic index (GI), such as whole grains and legumes, are absorbed more gradually, helping to maintain stable blood glucose levels and reduce insulin spikes (80). Bioactive compounds, such as polyphenols and flavonoids, found in functional foods enhance insulin sensitivity, which is crucial for efficient glucose

metabolism (7,79). The gut microbiota plays a key role in glucose regulation, and functional foods can positively influence gut health, thereby improving insulin function and reducing inflammation associated with diabetes (8). While functional foods show potential in managing blood glucose levels, their effectiveness may vary depending on individual responses and the specific characteristics of the foods consumed (81).

The role of functional foods in modulating inflammation and oxidative stress

Functional foods are rich in nutrients and bioactive compounds that help regulate inflammation and oxidative stress, contributing to overall health. Epidemiological studies indicate that diets high in fruits, vegetables, and fermented foods, such as the Mediterranean diet, are linked to lower inflammation levels and a decreased risk of chronic diseases (82). Certain bioactive compounds, such as polyphenols and flavonoids, can activate pathways like Nrf2, which controls responses to oxidative stress (83). These functional foods impact important key signaling pathways, including Nrf2 and NF- κ B, which are crucial for managing oxidative stress and inflammation (84).

Changes in gut microbiota and cellular signaling pathways

Dysbiosis, an imbalance in the microbiota, is linked to chronic inflammation and plays a role in conditions such as obesity and inflammatory bowel disease. Short-chain fatty acids (SCFAs) produced by the microbiota can help reduce the production of pro-inflammatory cytokines, thus regulating immune responses (85). The gut microbiota also influences important signaling pathways, such as the Toll-like receptor (TLR) pathway, which is crucial for immune system function. Dysbiosis can activate TLR signaling, leading to inflammatory processes associated with metabolic disorders (86). Targeting the gut microbiota with probiotics, prebiotics, and fecal microbiota transplantation presents new

therapeutic approaches for managing inflammatory diseases and metabolic disorders (87).

Table 1 outlines the main bioactive categories, their mechanisms of action, and the physiological effects relevant to diabetes management.

Conclusion

Research has shown that functional foods can improve glycemic control and insulin sensitivity, which are essential for managing diabetes (81). These foods have antioxidant and anti-inflammatory properties that can help reduce oxidative stress and inflammation linked to diabetes, potentially lowering the risk of complications (38). Modifying the gut microbiota through dietary changes has been found to impact glucose regulation,

human trials (88,89). Therefore, integrating functional foods into diabetes management should be done under professional supervision to ensure appropriate dietary modifications and prevent potential adverse effects. Future studies should prioritize robust trial designs and transparent reporting to strengthen the evidence base.

Acknowledgments

The author gratefully acknowledges the valuable insights gained from the scientific literature and the constructive feedback provided by peer reviewers, which have helped improve the clarity and rigor of this review.

Table 1. Mechanisms of action of functional foods in diabetes management

Bioactive category	Mechanistic pathways	Physiological effects
Dietary fibers	SCFA production, delayed glucose absorption	improved insulin sensitivity, glycemic control
Polyphenols	enzyme inhibition, AMPK activation	reduced postprandial glucose, anti-inflammatory effects
Antioxidants	ROS scavenging, Nrf2 activation	β -cell protection, reduced oxidative stress
Probiotics	microbiota modulation, butyrate production	reduced inflammation, enhanced metabolic function
Prebiotics	growth of beneficial bacteria, SCFA production	improved insulin sensitivity, reduced insulin resistance
Beneficial fatty acids	anti-inflammatory cytokine reduction, insulin signaling enhancement	lower insulin resistance, systemic inflammation reduction
Fermented foods	bioactive peptide and organic acid production	improved glycemic control, microbiota regulation
Plant compounds	enzyme inhibition, metabolic pathway regulation	reduced glucose absorption, improved insulin function

underscoring the importance of functional foods in supporting gut health and metabolic balance (8). However, the effectiveness of functional foods remains uncertain due to limitations in current research, such as small sample sizes, short study durations, and methodological differences. Individual factors like genetics and existing health conditions can significantly influence the impact of functional foods, making generalizations challenging (81). It's worth noting that much of the evidence supporting of functional foods comes from in vitro, animal, or preliminary

Funding

This work was conducted independently, without any external funding or institutional support.

Conflict of Interest

The author declares no conflicts of interest.

Authors' contributions

The author was responsible for the conception, literature review, critical analysis, manuscript drafting, and final approval of the version to be published.

References

- Nadhiya J, Vijayalakshmi MK, Showbharnikhaa S. A brief review on diabetes mellitus. *Journal of Pharma Insights and Research*. 2024;2(1):117-21.
- Pradhan D, Sahu PK, Purohit S, Ranajit SK, Acharya B, Sangam S, et al. Therapeutic interventions for diabetes mellitus-associated

- complications. *Current Diabetes Reviews*. 2025;21(8):e030524229631.
3. Karachaliou F, Simatos G, Simatou A. The challenges in the development of diabetes prevention and care models in low-income settings. *Frontiers in endocrinology*. 2020;11:518.
 4. Apini R, Annan RA, Apprey C, Asamoah-Boakye O. Predictors of glycaemic control among Ghanaian type 2 diabetic patients using diabetes self-management approach. *International Journal of Biochemistry Research & Review*. 2018; 23(2):1-18.
 5. das Chagas LA, de Sousa RM, Mattar R, Júnior JE, Júnior EA. Diet as a Risk Factor for the Development of Gestational Diabetes Mellitus: A Narrative Review [Internet]. *Public Health*. Intech Open 2025.; Available from: <http://dx.doi.org/10.5772/intechopen.1009119>.
 6. Mezouara M, Hbyaj K, Soufiani A, Mouniane Y, Chriqui A, Ramdane KB, et al. Potential of Functional Foods in Diabetes Management: A Review of *Coriandrum sativum*, *Ficus carica*, *Camellia sinensis*. *Advancements in Climate and Smart Environment Technology*. 2024;88-99.10.4018/979-8-3693-3807-0.ch008
 7. Li Sh. Application of Functional Food in Blood Glucose Regulation. *Highlights in Science, Engineering and Technology* 2024; 123: 194-199.
 8. Patel N, Dinesh S, Sharma S. From gut to glucose: a comprehensive review on functional foods and dietary interventions for diabetes management. *Current Diabetes Reviews*. 2024;20(5):23-44.
 9. Kayode AA, Adebodun GO, Fabunmi DI, Kayode OT. Role of functional foods in the management of diabetes mellitus: a concise review. *Bioactive Molecules and Pharmaceuticals*. 2023; 2(7): 29-41.
 10. Kaur J, Kaur K, Singh B, Singh A, Sharma S. Insights into the latest advances in low glycemic foods, their mechanism of action and health benefits. *Journal of Food Measurement and Characterization*. 2022;16(1):533-46.
 11. Silva FM, Kramer CK, de Almeida JC, Steemburgo T, Gross JL, Azevedo MJ. Fiber intake and glycemic control in patients with type 2 diabetes mellitus: a systematic review with meta-analysis of randomized controlled trials. *Nutrition reviews*. 2013;71(12):790-801.
 12. Chandalia M, Garg A, Lutjohann D, Von Bergmann K, Grundy SM, Brinkley LJ. Beneficial effects of high dietary fiber intake in patients with type 2 diabetes mellitus. *New England Journal of Medicine*. 2000;342(19):1392-8.
 13. Zhang R, Noronha JC, Khan TA, McGlynn N, Back S, Grant SM, et al. The effect of non-nutritive sweetened beverages on postprandial glycemic and endocrine responses: a systematic review and network meta-analysis. *Nutrients*. 2023;15(4):1050.
 14. Chen V, Zurbau A, Ahmed A, Khan TA, Au-Yeung F, Chiavaroli L, Mejia SB, et al. Effect of oats and oat β -glucan on glycemic control in diabetes: a systematic review and meta-analysis of randomized controlled trials. *BMJ Open Diabetes Research & Care*. 2022;10(5): e002784.
 15. Tessari P, Lante A. A multifunctional bread rich in beta glucans and low in starch improves metabolic control in type 2 diabetes: a controlled trial. *Nutrients*. 2017;9(3):297.
 16. Baroni I, Fabrizi D, Luciani M, Magon A, Conte G, De Angeli G, et al. Probiotics and synbiotics for glycemic control in diabetes: A systematic review and meta-analysis of randomized controlled trials. *Clinical Nutrition*. 2024;43(4):1041-61.
 17. Temple NJ. A rational definition for functional foods: A perspective. *Frontiers in nutrition*. 2022; 9:957516.
 18. Crowe KM, Francis C. Position of the academy of nutrition and dietetics: functional foods. *Journal of the Academy of Nutrition and Dietetics*. 2013;113(8):1096-103.
 19. Domínguez Díaz L, Fernández-Ruiz V, Cámara M. The frontier between nutrition and pharma: The international regulatory framework of functional foods, food supplements and nutraceuticals. *Critical reviews in food science and nutrition*. 2020;60(10):1738-46.
 20. Moraes FP. Alimentos funcionais e nutracêuticos: definições, legislação e benefícios à saúde. *Revista eletrônica de farmácia*. 2006;3(2). <https://doi.org/10.5216/ref.v3i2.2082>
 21. Phillips MM, Rimmer CA. Functional foods and dietary supplements. *Analytical and bioanalytical chemistry*. 2013;405(13):4323-4.
 22. Martirosyan D, Ekblad M. Functional foods classification system: exemplifying through analysis of bioactive compounds. *Functional Food Science-Online* ISSN: 2767-3146. 2022;2(4):94-123.
 23. Thompson AK, Moughan PJ. Innovation in the foods industry: Functional foods. *Innovation*. 2008;10(1):61-73.
 24. Helkar PB, Sahoo AK, Patil NJ. Review: Food industry by-products used as a functional food ingredients. *Int. J. Waste Resour*. 2016;6(3):1-6.
 25. Ospina-Maldonado S, Martín-Gómez H, Cardoso-Ugarte GA. From waste to wellness: a review on the harness of food industry by-products for sustainable functional food production. *International Journal of Food Science and Technology*. 2024;59(11):8680-92.
 26. Granato D, Barba FJ, Bursac Kovačević D, Lorenzo JM, Cruz AG, Putnik P. Functional foods: Product development, technological trends, efficacy testing, and safety. *Annual review of food science and technology*. 2020;11(1):93-118.

27. Sivamaruthi BS, Kesika P, Prasanth MI, Chaiyasut C. A mini review on antidiabetic properties of fermented foods. *Nutrients*. 2018;10(12):1973.
28. Wang W, Yang W, Dai Y, Liu J, Chen ZY. Production of food-derived bioactive peptides with potential application in the management of diabetes and obesity: A review. *Journal of Agricultural and Food Chemistry*. 2023;71(15):5917-43.
29. Asryan Vr. Biotechnology in the production of functional food products. *Collection of Scientific Papers*. 2024.
30. Barma NR, Hossen NM. Modern technologies in dieting and diabetic management. *Global Journal of Engineering & Technology Advances* 2023; 16(2):227-34.
31. Giacco R, De Giulio B, Vitale M, Cozzolino R. Functional Foods: Can Food Technology Help in the Prevention and Treatment of Diabetes?. *Food and Nutrition Sciences*. 2013;4(8):827-37.
32. Alba JB, Guambuguete CE, Hachi AL, Jácome JEQC. Functional Foods: General Analysis and Development in the Food Industry. *International Journal of Current Microbiology & Applied Sciences* 2021; 10(12): 220-230. <https://www.ijcmas.com/10-12-2021/Jimmy%20Bladimir%20Alba,%20et%20al.pdf>
33. Mogoş T, Dondoi C, Iacobini AE. A review of dietary fiber in the diabetic diet. *Romanian Journal of Diabetes Nutrition and Metabolic Diseases*. 2017;24(2):161-4.
34. Mazhar M, Zhu Y, Qin L. The interplay of dietary fibers and intestinal microbiota affects type 2 diabetes by generating short-chain fatty acids. *Foods*. 2023;12(5):1023.
35. Lattimer JM, Haub MD. Effects of dietary fiber and its components on metabolic health. *Nutrients*. 2010;2(12):1266-89.
36. Reynolds AN, Akerman AP, Mann J. Dietary fibre and whole grains in diabetes management: Systematic review and meta-analyses. *PLoS medicine*. 2020;17(3):e1003053.
37. Raina J, Firdous A, Singh G, Kumar R, Kaur C. Role of polyphenols in the management of diabetic complications. *Phytomedicine*. 2024 Jan 1;122:155155.
38. Aryaeian N, Sedehi SK, Arablou T. Polyphenols and their effects on diabetes management: A review. *Medical journal of the Islamic Republic of Iran*. 2017;31:134.
39. Barik SK, Sengupta S, Arya R, Kumar S, Kim JJ, Chaurasia R. Dietary Polyphenols as Potential Therapeutic Agents in Type 2 Diabetes Management: Advances and Opportunities. *Advances in Nutrition*. 2025;16(1):100346.
40. Serina JJ, Castilho PC. Using polyphenols as a relevant therapy to diabetes and its complications, a review. *Critical Reviews in Food Science and Nutrition*. 2022;62(30):8355-87.
41. Guasch-Ferré M, Merino J, Sun Q, Fitó M, Salas-Salvadó J. Dietary polyphenols, Mediterranean diet, prediabetes, and type 2 diabetes: a narrative review of the evidence. *Oxidative medicine and cellular longevity*. 2017;2017(1):6723931.
42. David BG. Polyphenols and Insulin Sensitivity: A Pathophysiological Perspective. *Research Output Journal of Public Health & Medicine* 2025; 5(3):12-15.
43. Kim Y, Keogh JB, Clifton PM. Polyphenols and glycemic control. *Nutrients*. 2016;8(1):17.
44. Xu D, Fu L, Pan D, Chu Y, Feng M, Lu Y, et al. Role of probiotics/synbiotic supplementation in glycemic control: A critical umbrella review of meta-analyses of randomized controlled trials. *Critical reviews in food science and nutrition*. 2024;64(6):1467-85.
45. Singh H, Singh R, Singh A, Singh H, Singh G, Kaur S, et al. Role of oxidative stress in diabetes-induced complications and their management with antioxidants. *Archives of Physiology and Biochemistry*. 2024;130(6):616-41.
46. Tumilaar SG, Hardianto A, Dohi H, Kurnia D. A comprehensive review of free radicals, oxidative stress, and antioxidants: Overview, clinical applications, global perspectives, future directions, and mechanisms of antioxidant activity of flavonoid compounds. *Journal of Chemistry*. 2024;2024(1):5594386.
47. Oh YS, Jun HS. Role of bioactive food components in diabetes prevention: Effects on Beta-cell function and preservation. *Nutrition and Metabolic Insights*. 2014:NMI-S13589.
48. Zhong O, Hu J, Wang J, Tan Y, Hu L, Lei X. Antioxidant for treatment of diabetic complications: A meta-analysis and systematic review. *Journal of Biochemical and Molecular Toxicology*. 2022;36(6):e23038.
49. Banaszak M, Górna I, Woźniak D, Przysławski J, Drzymała-Czyż S. The impact of curcumin, resveratrol, and cinnamon on modulating oxidative stress and antioxidant activity in type 2 diabetes: moving beyond an anti-hyperglycaemic evaluation. *Antioxidants*. 2024;13(5):510.
50. Choi SW, Ho CK. Antioxidant properties of drugs used in Type 2 diabetes management: could they contribute to, confound or conceal effects of antioxidant therapy?. *Redox Report*. 2018 ;23(1):1-24.
51. Sanz-González SM, García-Medina JJ, Zanón-Moreno V, López-Gálvez MI, Galarreta-Mira D, Duarte L, et al. Clinical and molecular-genetic insights into the role of oxidative stress in diabetic retinopathy: antioxidant strategies and future avenues. *Antioxidants*. 2020;9(11):1101.

52. Sun L, Ni C, Zhao J, Wang G, Chen W. Probiotics, bioactive compounds and dietary patterns for the effective management of hyperuricemia: a review. *Critical Reviews in Food Science and Nutrition*. 2024;64(7):2016-31.
53. Li S, Liu Z, Zhang Q, Su D, Wang P, Li Y, et al. The antidiabetic potential of probiotics: a review. *Nutrients*. 2024;16(15):2494.
54. Sooragonda BG, Thakur P, Moganti R, Garg A, Garg UK. Systematic Review of the Effectiveness of Common Probiotic Strains: *Streptococcus faecalis*, *Clostridium butyricum*, *Bacillus mesentericus*, and *Lactobacillus sporogenes* in the Management of Diabetes. *Journal of Diabetology*. 2025;16(1):29-36.
55. Sreepathi N, Jayanthi MK, Chandra SJ, Bajpe SN, Ramu R. Probiotic intervention in the treatment of diabetes mellitus: a review. *Journal of Pure and Applied Microbiology* 2022; 6:1519-29.
56. Ayesha IE, Monson NR, Klair N, Patel U, Saxena A, Patel D, et al. Probiotics and their role in the management of type 2 diabetes mellitus (Short-term versus long-term effect): A systematic review and meta-analysis. *Cureus*. 2023;15(10):e46741.
57. Okburan G, Bas M, Ogmen S. A randomized double-blind controlled clinical trial demonstrating efficacy of different probiotic strains on serum lipids and glycemic biomarker. *Nutrición Hospitalaria*. 2024;41(4):793-803.
58. Osadebe PO, Odoh EU, Uzor PF. The search for new hypoglycemic agents from plants. *Afr J Pharm Pharmacol*. 2014;8(11):292-303.
59. Iatcu OC, Hamamah S, Covasa M. Harnessing prebiotics to improve type 2 diabetes outcomes. *Nutrients*. 2024;16(20):3447.
60. Baghel K, Khan A, Kango N. Role of synbiotics (prebiotics and probiotics) as dietary supplements in type 2 diabetes mellitus induced health complications. *Journal of Dietary Supplements*. 2024;21(5):677-708.
61. Kumar A, Prajapati B, Pramanik J. Prebiotics as adjunctive therapy in diabetes: a review of prebiotics in diabetes. *Current Nutraceuticals*. 2022;3(3):e180822207618.
62. Sharma BR, Jaiswal S, Ravindra PV. Modulation of gut microbiota by bioactive compounds for prevention and management of type 2 diabetes. *Biomedicine & Pharmacotherapy*. 2022;152:113148.
63. Das UN. Is there a role for bioactive lipids in the pathobiology of diabetes mellitus?. *Frontiers in endocrinology*. 2017;8:182.
64. Shetty SS, Kumari S. Fatty acids and their role in type-2 diabetes. *Experimental and therapeutic medicine*. 2021;22(1):1-6.
65. Rice Bradley BH. Dietary fat and risk for type 2 diabetes: a review of recent research. *Current nutrition reports*. 2018;7(4):214-26.
66. Sivri D, Akdevelioğlu Y. Effect of fatty acids on glucose metabolism and type 2 diabetes. *Nutrition Reviews*. 2025;83(5):897-907.
67. Melini F, Melini V, Luziatelli F, Ficca AG, Ruzzi M. Health-promoting components in fermented foods: An up-to-date systematic review. *Nutrients*. 2019;11(5):1189.
68. Deveci G, Çelik E, Ağagündüz D, Bartkiene E, Rocha JM, Özogul F. Certain fermented foods and their possible health effects with a focus on bioactive compounds and microorganisms. *Fermentation*. 2023;9(11):923.
69. Kurian C, Mathur A, Paari K. Review on dietary factors in fermented foods and their efficacy in disease management. *Current Nutrition & Food Science*. 2022;18(2):144-65.
70. Liu L, Li G, Cui L, Cai R, Yuan Y, Gao Z, et al. The health benefits of fermented fruits and vegetables and their underlying mechanisms. *Comprehensive reviews in food science and food safety*. 2024;23(6):e70072.
71. de Almeida Souza C, de Oliveira ÍA, de Oliveira Rolim VA, Bogsan CS. Traditional fermented foods as an adjuvant treatment to diabetes. *Current Geriatrics Reports*. 2020;9(4):242-50.
72. Ostadrahimi A, Taghizadeh A, Mobasseri M, Farrin N, Payahoo L, Gheshlaghi ZB, et al. Effect of probiotic fermented milk (kefir) on glycemic control and lipid profile in type 2 diabetic patients: a randomized double-blind placebo-controlled clinical trial. *Iranian journal of public health*. 2015;44(2):228.
73. Ahmad R, Oli AN, Etando A, Sharma P, Sinha S, Chowdhury K, et al. Lactic acid bacteria fermented foods: Impact on immune system and consequences over type 2 diabetes mellitus. *Journal of Applied Pharmaceutical Science*. 2023;13(6):018-56.
74. Onukwuli CO, Izuchukwu E, Paul-Chima O. Exploring phytochemicals for diabetes management: Mechanisms, efficacy, and future directions. *SCIENCES (NIJ RMS)*. 2024;5(2).
75. El-Nashar HA, Mostafa NM, El-Shazly M, Eldahshan OA. The role of plant-derived compounds in managing diabetes mellitus: a review of literature from 2014 to 2019. *Current medicinal chemistry*. 2021;28(23):4694-730.
76. Mustafa AM, Dawod OY, Mohammed RA, Alfaki AM, Ahmed SOM, Abdelrahman MH, et al. Exploring Sudanese Medicinal Plants for their Antidiabetic and Hypolipidemic Benefits: A Review. *South Eastern European Journal of Public Health* 2025: 4095–4111. <https://doi.org/10.70135/seejph.vi.4768>
77. Salleh NH, Zulkipli IN, Mohd Yasin H, Ja'afar F, Ahmad N, Wan Ahmad WA, et al. Systematic review of medicinal plants used for treatment of diabetes in human clinical trials: An ASEAN

- perspective. *Evidence-Based Complementary and Alternative Medicine*. 2021;2021(1):5570939.
78. Solnier J, Chang C, Pizzorno J. Consideration for flavonoid-containing dietary supplements to tackle deficiency and optimize health. *International journal of molecular sciences*. 2023;24(10):8663.
79. Mirmiran P, Bahadoran Z, Azizi F. Functional foods-based diet as a novel dietary approach for management of type 2 diabetes and its complications: A review. *World journal of diabetes*. 2014;5(3):267.
80. Murarka S, Singh SP. Review on low glycaemic index functional food products. *Pharma Innovation Journal* 2023; 12:624-32.
81. Hamasaki H. Functional foods for type 2 diabetes. *AIMS Medical Science*. 2016;3(3):278-97.
82. Gok I. Bioactive compounds modulating inflammation and oxidative stress in some traditional functional foods and beverages. In *Current advances for development of functional foods modulating inflammation and oxidative stress*. Academic Press .2022: 1-16.
83. Tkaczenko H, Kurhaluk N. Antioxidant-rich functional foods and exercise: unlocking metabolic health through Nrf2 and related pathways. *International Journal of Molecular Sciences*. 2025;26(3):1098.
84. Martins-Gomes C, Nunes FM, Silva AM. Natural products as dietary agents for the prevention and mitigation of oxidative damage and inflammation in the intestinal barrier. *Antioxidants*. 2024;13(1):65.
85. Nigam M, Devi K, Coutinho HD, Mishra AP. Exploration of gut microbiome and inflammation: A review on key signalling pathways. *Cellular Signalling*. 2024;118:111140.
86. Chen L, Zhang L, Hua H, Liu L, Mao Y, Wang R. Interactions between toll-like receptors signaling pathway and gut microbiota in host homeostasis. *Immunity, Inflammation and Disease*. 2024;12(7):e1356.
87. Shi Y, Chen Z, Fang T, Chen X, Deng Y, Qin H, Lian M, Shen J, Zong Y, Chu H, Hoebinger C. Gut microbiota in treating inflammatory digestive diseases: Current challenges and therapeutic opportunities. *iMeta*. 2025;4(1):e265..
88. Matsinkou Soh R, Ngaha Damndja W, Njintang Yanou N. Functional foods of sub-Saharan Africa and their implications in the management of type 2 diabetes: A review. *Food Science & Nutrition*. 2024;12(1):24-34.
89. Lam HN, Lin SP, Nguyen DH, Chen CM, Su CT, Fang TC, et al. Integrative Roles of Functional Foods, Microbiotics, Nutrigenetics, and Nutrigenomics in Managing Type 2 Diabetes and Obesity. *Nutrients*. 2025;17(4):608.